

Rabbit HMGB1 Polyclonal Antibody | anti-HMGB1 antibody

(KO Validated) HMGB1 Polyclonal Antibody

CATALOG # AAA178723	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

HMGB1; Polyclonal Antibody; (KO Validated) HMGB1 Polyclonal Antibody; HMG-1; HMG1; HMG3; SBP-1; anti-HMGB1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.01% thiomersal, 50% glycerol, pH 7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-HMGB1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human HMGB1
Research Area	Cancer, Epigenetics and Nuclear Signaling
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse heart using HMGB1 Polyclonal antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human breast cancer using HMGB1 Polyclonal antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat brain using HMGB1 Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of various lysates using HMGB1 Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-HMGB1 antibody	This gene encodes a protein that belongs to the High Mobility Group-box superfamily. The encoded non-histone, nuclear DNA-binding protein regulates transcription, and is involved in organization of DNA. This protein plays a role in several cellular processes, including inflammation, cell differentiation and tumor cell migration. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants that encode the same protein.
Product Categories/Family for anti-HMGB1 antibody	Antibodies ; Primary Antibodies ; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504425
NCBI GeneID	3146
NCBI Accession #	NP_002119.1
NCBI GenBank Nucleotide #	NM_002128.4
UniProt Accession #	P09429 ; Q14321 ; Q5T7C3 ; Q6IBE1 ; A5D8W9
Molecular Weight	24,894 Da
NCBI Official Full Name	high mobility group protein B1
NCBI Official Synonym Full Names	high mobility group box 1
NCBI Official Symbol	HMGB1
NCBI Official Synonym Symbols	HMG1; HMG3; SBP-1
NCBI Protein Information	high mobility group protein B1; HMG-1; Amphoterin; high-mobility group box 1; high mobility group protein 1; Sulfoglucuronyl carbohydrate binding protein; high-mobility group (nonhistone chromosomal) protein 1
UniProt Protein Name	High mobility group protein B1
UniProt Gene Name	HMGB1
UniProt Synonym Gene Names	HMG1; HMG-1
UniProt Entry Name	HMGB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

