

Rabbit PDCD1LG2 Polyclonal Antibody | anti-PDCD1LG2 antibody

PDCD1LG2 Polyclonal Antibody

CATALOG # AAA179135	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PDCD1LG2; Polyclonal Antibody; PDCD1LG2 Polyclonal Antibody; B7DC; Btdc; CD273; PD-L2; PDCD1L2; PDL2; bA574F11.2; anti-PDCD1LG2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-PDCD1LG2 antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthetic peptide of human PDCD1LG2
Research Area	Immunology
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of NIH/3T3 cells using PDCD1LG2 Polyclonal Antibody at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.



IF (Immunofluorescence)

Immunofluorescence analysis of HepG2 cells using PDCD1LG2 Polyclonal Antibody at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.



WB (Western Blot)
 Western blot analysis of extracts of various cell lines using PD-CD1LG2 Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-PD-CD1LG2 antibody	Programmed Death Ligand 2 (PD-L2), also known as B7-DC and butyrophilin-like protein, is member of the B7 family of proteins that provide signals for regulating T-cell activation and tolerance. PD-L2 is expressed on dendritic cells, subsets of activated CD4+ and CD8+ T cells, and memory B cells that differentiate into plasma cells. At inflammatory sites such as rheumatoid arthritis, allergen exposure, and virus infection, PD-L2 is up-regulated on synoviocytes, infiltrating macrophages, dendritic cells, and airway epithelial cells. PD-L2, along with B7-H1/PD-L1, binds to T cell PD-1 where it promotes IFN-gamma production and CD40 Ligand up-regulation while inhibiting IL-4 production. In addition, PD-L2 binds to RGM-B on macrophages and alveolar epithelial cells, supporting respiratory immune tolerance. In asthma, PD-L2 suppresses IL-5 and IL-13 production, promotes IL-12 production by dendritic cells, and supports allergen-induced airway hyper-responsiveness and mucus production.
Product Categories/Family for anti-PD-CD1LG2 antibody	Antibodies; Primary Antibodies; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	190014605
NCBI GeneID	80380
NCBI Accession #	NP_079515.2
NCBI GenBank Nucleotide #	NM_025239.3
UniProt Accession #	Q9BQ51; Q14CN8; Q5T7Z6; Q6JXL8; Q6JXL9
Molecular Weight	20,745 Da
NCBI Official Full Name	programmed cell death 1 ligand 2
NCBI Official Synonym Full Names	programmed cell death 1 ligand 2
NCBI Official Symbol	PDCD1LG2
NCBI Official Synonym Symbols	B7DC; Btdc; PDL2; CD273; PD-L2; PDCD1L2; bA574F11.2
NCBI Protein Information	programmed cell death 1 ligand 2
UniProt Protein Name	Programmed cell death 1 ligand 2
UniProt Gene Name	PDCD1LG2
UniProt Synonym Gene Names	B7DC; CD273; PDCD1L2; PDL2; PD-1 ligand 2; PD-L2; PDCD1 ligand 2; Programmed death ligand 2; B7-DC
UniProt Entry Name	PD1L2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/pdcd1lg2-antibody-immunofluorescence-western-blot-elisa/179135>