

Rabbit anti-Human CSTB Polyclonal Antibody | anti-CSTB antibody

CSTB Polyclonal Antibody

CATALOG # AAA179322	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CSTB; Polyclonal Antibody; CSTB Polyclonal Antibody; CPI-B; CST6; EPM1; EPM1A; PME; STFB; ULD; cystatin-B; anti-CSTB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CSTB antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human CSTB
Research Area	Cancer, Cell Biology
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon carcinoma using CSTB Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human thyroid cancer using CSTB Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of various cell lines using CSTB Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-CSTB antibody

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and kininogens. This gene encodes a stefin that functions as an intracellular thiol protease inhibitor. The protein is able to form a dimer stabilized by noncovalent forces, inhibiting papain and cathepsins I, h and b. The protein is thought to play a role in protecting against the proteases leaking from lysosomes. Evidence indicates that mutations in this gene are responsible for the primary defects in patients with progressive myoclonic epilepsy (EPM1). One type of mutation responsible for EPM1 is the expansion in the promoter region of this gene of a CCCC GCCCGCG repeat from 2-3 copies to 30-78 copies.

Product Categories/Family for anti-CSTB antibody

[Antibodies](#); [Primary Antibodies](#); [Polyclonal Antibody](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503117
NCBI GeneID	1476
NCBI Accession #	NP_000091.1
NCBI GenBank Nucleotide #	NM_000100.3
UniProt Accession #	P04080
Molecular Weight	11,140 Da
NCBI Official Full Name	cystatin-B
NCBI Official Synonym Full Names	cystatin B (stefin B)
NCBI Official Symbol	CSTB
NCBI Official Synonym Symbols	PME; ULD; CST6; EPM1; STFB; EPM1A
NCBI Protein Information	cystatin-B; CPI-B; liver thiol proteinase inhibitor
UniProt Protein Name	Cystatin-B
UniProt Gene Name	CSTB
UniProt Synonym Gene Names	CST6; STFB
UniProt Entry Name	CYTB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cstb-antibody-immunohistochemistry-western-blot-elisa/179322>