

Rabbit M6PR Polyclonal Antibody | anti-M6PR antibody

M6PR Polyclonal Antibody

CATALOG # AAA179329	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

M6PR; Polyclonal Antibody; M6PR Polyclonal Antibody; CD-M6PR; CD-MPR; MPR 46; MPR-46; MPR46; SMPR; cation-dependent mannose-6-phosphate receptor; anti-M6PR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-M6PR antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human M6PR
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse heart using M6PR Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using M6PR Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver damage using M6PR Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of mouse kidney using M6PR Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-M6PR antibody	This gene encodes a member of the P-type lectin family. P-type lectins play a critical role in lysosome function through the specific transport of mannose-6-phosphate-containing acid hydrolases from the Golgi complex to lysosomes. The encoded protein functions as a homodimer and requires divalent cations for ligand binding. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. A pseudogene of this gene is located on the long arm of chromosome X.
Product Categories/Family for anti-M6PR antibody	Antibodies ; Primary Antibodies ; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505061
NCBI GeneID	4074
NCBI Accession #	NP_002346
NCBI GenBank Nucleotide #	NM_002355.3
UniProt Accession #	P20645 ; A8K528 ; D3DUV5
Molecular Weight	30,993 Da
NCBI Official Full Name	cation-dependent mannose-6-phosphate receptor isoform 1
NCBI Official Synonym Full Names	mannose-6-phosphate receptor, cation dependent
NCBI Official Symbol	M6PR
NCBI Official Synonym Symbols	SMPR; MPR46; CD-MPR; MPR 46; MPR-46; CD-M6PR
NCBI Protein Information	cation-dependent mannose-6-phosphate receptor
UniProt Protein Name	Cation-dependent mannose-6-phosphate receptor
UniProt Gene Name	M6PR
UniProt Synonym Gene Names	MPR46; MPRD; CD Man-6-P receptor; CD-MPR; MPR 46

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/m6pr-antibody-immunohistochemistry-western-blot-elisa/179329>