

Rabbit Cyclin E1 Polyclonal Antibody | anti-CCNE1 antibody

Cyclin E1 Polyclonal Antibody

CATALOG # AAA179491	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Cyclin E1; Polyclonal Antibody; Cyclin E1 Polyclonal Antibody; CCNE1; CCNE; pCCNE1; cyclin E1; anti-CCNE1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CCNE1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human Cyclin E1
Research Area	Cancer, Cell Biology, Epigenetics and Nuclear Signaling
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse lung using Cyclin E1 Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of various cell lines using Cyclin E1 Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-CCNE1 antibody	The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB.
Product Categories/Family for anti-CCNE1 antibody	Antibodies; Primary Antibodies; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	17318559
NCBI GeneID	898
NCBI Accession #	NP_001229.1
NCBI GenBank Nucleotide #	NM_001238.2
UniProt Accession #	P24864; Q14091; Q8NFG1; Q92501; Q9UD21; A8K684
Molecular Weight	47,077 Da
NCBI Official Full Name	G1/S-specific cyclin-E1
NCBI Official Synonym Full Names	cyclin E1
NCBI Official Symbol	CCNE1
NCBI Official Synonym Symbols	CCNE
NCBI Protein Information	G1/S-specific cyclin-E1; cyclin Es; cyclin Et
UniProt Protein Name	G1/S-specific cyclin-E1
UniProt Gene Name	CCNE1
UniProt Synonym Gene Names	CCNE
UniProt Entry Name	CCNE1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cyclin-e1-antibody-immunohistochemistry-western-blot-elisa/179491>