

Rabbit anti-Human CK-18 Polyclonal Antibody | anti-CK-18 antibody

CK-18 Polyclonal Antibody

CATALOG # AAA179762	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CK-18; Polyclonal Antibody; CK-18 Polyclonal Antibody; Cell proliferation inducing gene 46 protein; Cell proliferation inducing protein 46; Cell proliferation-inducing gene 46 protein; CK 18; CK18; CYK 18; CYK18; Cytokeratin 18; Cytokeratin endo B; Cytokeratin-18; K 18; K18; K1C18; KA18; Keratin 18; type I; Keratin D; keratin; type I cytoskeletal 18; Keratin-18; Krt18; anti-CK-18 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purification
Form/Format	PBS with 0.02% sodium azide, 1% protective protein and 50% glycerol, pH 7.4
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CK-18 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant Human KRT18 protein expressed by E coli
Preparation and Storage	Store at -20 degree C Valid for 12 months. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon cancer tissue using AGR1 Polyclonal Antibody at dilution of 1:2900




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human lung cancer tissue using AGR1 Polyclonal Antibody at dilution of 1:2900



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using AGR1 Polyclonal Antibody at dilution of 1:2900



WB (Western Blot)

Western blot with AGR1 Polyclonal antibody at dilution of 1:5000. lane 1: Hela whole cell lysate, lane 2: A431 whole cell lysate

ADDITIONAL INFORMATION

Related Product Information for anti-CK-18 antibody	Cytokeratin 18 (CK18) has a molecular weight of 45 KDa, which is a low molecular weight type A cytokeratin. It is a marker of simple epithelium, including glandular epithelium. The stratified squamous epithelium is generally negative. CK18 is mainly used in the identification of adenocarcinoma.
Product Categories/Family for anti-CK-18 antibody	Polyclonal Antibodies ; Primary Antibodies ; Polyclonal Antibodies ; Cancer ; Signal Transduction ; Tags and Cell Markers

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557888
NCBI GeneID	3875
NCBI Accession #	NP_000215.1
NCBI GenBank Nucleotide #	NM_000224.2
UniProt Accession #	P05783; Q53G38; Q5U0N8; Q9BW26
Molecular Weight	48,058 Da
NCBI Official Full Name	keratin, type I cytoskeletal 18
NCBI Official Synonym Full Names	keratin 18
NCBI Official Symbol	KRT18
NCBI Official Synonym Symbols	K18; CYK18
NCBI Protein Information	keratin, type I cytoskeletal 18; CK-18; keratin-18; cytokeratin 18; cytokeratin-18; cell proliferation-inducing protein 46; cell proliferation-inducing gene 46 protein
UniProt Protein Name	Keratin, type I cytoskeletal 18
UniProt Gene Name	KRT18
UniProt Synonym Gene Names	CYK18; CK-18; K18
UniProt Entry Name	K1C18_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ck-18-antibody-immunohistochemistry-western-blot-elisa/179762>