

Dog Adrenomedullin (ADM) ELISA Kit | ADM elisa kit

Dog Adrenomedullin (ADM) ELISA Kit

CATALOG #
AAA181877

REACTIVITY
Dog


PRODUCT SYNONYMS

Adrenomedullin (ADM); Dog Adrenomedullin (ADM) ELISA Kit; AM; PAMP; Proadrenomedullin N-20 terminal peptide; ProAM N-terminal 20 peptide; ADM elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Dog
Specificity	This assay has high sensitivity and excellent specificity for detection of Adrenomedullin (ADM). No significant cross-reactivity or interference between Adrenomedullin (ADM) and analogues was observed.
Sequence Length	185
Assay Type	Competitive Inhibition
Samples	Serum, Plasma and Other Biological Fluids
Detection Range	12.35-1,000pg/mL
Sensitivity	< 4.42pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Adrenomedullin (ADM) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Adrenomedullin (ADM) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ADM elisa kit	This assay employs the competitive inhibition enzyme immunoassay technique. A monoclonal antibody specific to Adrenomedullin (ADM) has been pre-coated onto a microplate. A competitive inhibition reaction is launched between biotin labeled Adrenomedullin (ADM) and unlabeled Adrenomedullin (ADM) (Standards or samples) with the pre-coated antibody specific to Adrenomedullin (ADM). After incubation the unbound conjugate is washed off. Next, avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. The amount of bound HRP conjugate is reverse proportional to the concentration of Adrenomedullin (ADM) in the sample. After addition of the substrate solution, the intensity of color developed is reverse proportional to the concentration of Adrenomedullin (ADM) in the sample.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	765330
NCBI GeneID	133
UniProt Accession #	P35318; Q6FGW2; B2R793; D3DQV3
Molecular Weight	20,420 Da
NCBI Official Full Name	adrenomedullin
NCBI Official Synonym Full Names	adrenomedullin
NCBI Official Symbol	ADM
NCBI Official Synonym Symbols	AM; PAMP
NCBI Protein Information	ADM
UniProt Protein Name	ADM
UniProt Gene Name	ADM
UniProt Synonym Gene Names	AM; AM; PAMP; ProAM-N20

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/dog-elisa-kits/adrenomedullin-adm/181877>