

Dog Interleukin 3 (IL3) ELISA Kit | IL3 elisa kit

Dog Interleukin 3 (IL3) ELISA Kit

CATALOG #
AAA182732

REACTIVITY
Dog


PRODUCT SYNONYMS

Interleukin 3 (IL3); Dog Interleukin 3 (IL3) ELISA Kit; MCGF; MULTI-CSF; MCSF; M-CSF; Multiple; Multilineage-Colony-Stimulating Factor; Hematopoietic Growth Factor; P-Cell Stimulating Factor; Mast-Cell Growth Factor; IL3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Dog
Specificity	This assay has high sensitivity and excellent specificity for detection of Interleukin 3 (IL3). No significant cross-reactivity or interference between Interleukin 3 (IL3) and analogues was observed.
Sequence Length	152
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma and Other Biological Fluids
Detection Range	15.62-1000pg/mL
Sensitivity	< 6.5pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Interleukin 3 (IL3) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Interleukin 3 (IL3) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL3 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Interleukin 3 (IL3). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Interleukin 3 (IL3). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Interleukin 3 (IL3), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Interleukin 3 (IL3) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	28416915
NCBI GeneID	3562
NCBI Accession #	NP_000579.2
NCBI GenBank Nucleotide #	NM_000588.3
UniProt Accession #	P08700; Q6GS87
Molecular Weight	17,233 Da
NCBI Official Full Name	interleukin-3
NCBI Official Synonym Full Names	interleukin 3
NCBI Official Symbol	IL3
NCBI Official Synonym Symbols	IL-3; MCGF; MULTI-CSF
NCBI Protein Information	interleukin-3
UniProt Protein Name	Interleukin-3
UniProt Gene Name	IL3
UniProt Synonym Gene Names	IL-3; MCGF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.