

Porcine Monocyte Chemotactic Protein 2 (MCP2) ELISA Kit | MCP2 elisa kit

Pig Monocyte Chemotactic Protein 2 (MCP2) ELISA Kit

CATALOG #
AAA182805

REACTIVITY
Porcine


PRODUCT SYNONYMS

Monocyte Chemotactic Protein 2 (MCP2); Pig Monocyte Chemotactic Protein 2 (MCP2) ELISA Kit; CCL8; CCL-8; SCYA8; MCP-2; HC14; Chemokine C-C-Motif Ligand 8; Small Inducible Cytokine Subfamily A(Cys-Cys)Member 8; Monocyte Chemoattractant Protein 2; MCP2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Specificity	This assay has high sensitivity and excellent specificity for detection of MCP2. No significant cross-reactivity or interference between MCP2 and analogues was observed.
Sequence Length	99
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	15.6-1,000pg/mL
Sensitivity	< 6.3pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level MCP2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level MCP2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MCP2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of MCP2 in porcine serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to MCP2. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to MCP2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MCP2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of MCP2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
References	Specialized Proresolving Mediators Facilitate the Immunomodulation of the Periodontal Ligament Stem Cells; Frontiers in Dental Medicine; 2021 Volume 2; Ning Yu; Department of Applied Oral Sciences, The Forsyth Institute, Cambridge, MA, United States

NCBI AND UNIPROT INFORMATION

NCBI GI #	2916796
UniProt Accession #	P80075; P78388; A0AV77
Molecular Weight	11,246 Da
NCBI Official Full Name	monocyte chemotactic protein-2
UniProt Protein Name	C-C motif chemokine 8
UniProt Gene Name	CCL8
UniProt Synonym Gene Names	MCP2; SCYA10; SCYA8; MCP-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/porcine-elisa-kits/monocyte-chemotactic-protein-2-mcp2/182805>