

Rat Stromal Cell Derived Factor 1 (SDF1) ELISA Kit | SDF1 elisa kit

Rat Stromal Cell Derived Factor 1 (SDF1) ELISA Kit

CATALOG #
AAA182939

REACTIVITY
Rat


PRODUCT SYNONYMS

Stromal Cell Derived Factor 1 (SDF1); Rat Stromal Cell Derived Factor 1 (SDF1) ELISA Kit; CXCL12; PBSF; P-BSF; SCYB12; SDF1-A; SDF1-B; SDF1A; SDF1B; TLSF-A; TLSF-B; TPAR1; Pre-B Cell Growth-Stimulating Factor; Chemokine C-X-C-Motif Ligand 12; SDF1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of SDF1. No significant cross-reactivity or interference between SDF1 and analogues was observed.
Sequence Length	89
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	< 0.055ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SDF1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SDF1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SDF1 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of SDF1 in rat serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to SDF1. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to SDF1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SDF1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of SDF1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	40316924
NCBI GeneID	6387
NCBI Accession #	NP_954637.1
NCBI GenBank Nucleotide #	NM_199168.3
UniProt Accession #	P48061; Q2L985; Q2L986; Q2L988; Q5IT36; Q6ICW0; Q9H554; B2R4G0; E7EVL0; H7BYN8
Molecular Weight	11,004 Da
NCBI Official Full Name	stromal cell-derived factor 1 isoform alpha
NCBI Official Synonym Full Names	C-X-C motif chemokine ligand 12
NCBI Official Symbol	CXCL12
NCBI Official Synonym Symbols	IRH; PBSF; SDF1; TLSF; TPAR1; SCYB12
NCBI Protein Information	stromal cell-derived factor 1
UniProt Protein Name	Stromal cell-derived factor 1
UniProt Gene Name	CXCL12
UniProt Synonym Gene Names	SDF1; SDF1A; SDF1B; SDF-1; hSDF-1; IRH; hIRH; PBSF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/stromal-cell-derived-factor-1-sdf1/182939>