

Human Tissue Polypeptide Antigen (TPA) ELISA Kit | TPA elisa kit

Human Tissue Polypeptide Antigen (TPA) ELISA Kit

CATALOG #
AAA183059

REACTIVITY
Human


PRODUCT SYNONYMS

Tissue Polypeptide Antigen (TPA); Human Tissue Polypeptide Antigen (TPA) ELISA Kit; TPA elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Tissue Polypeptide Antigen (TPA). No significant cross-reactivity or interference between Tissue Polypeptide Antigen (TPA) and analogues was observed.
Sequence Length	19
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma and Other Biological Fluids
Detection Range	78.12-5000pg/mL
Sensitivity	< 29pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Tissue Polypeptide Antigen (TPA) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Tissue Polypeptide Antigen (TPA) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TPA elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Tissue Polypeptide Antigen (TPA). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Tissue Polypeptide Antigen (TPA). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Tissue Polypeptide Antigen (TPA), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Tissue Polypeptide Antigen (TPA) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	2285954
NCBI GeneID	5327
UniProt Accession #	Q6LBF5; P00750
Molecular Weight	2,342 Da
NCBI Official Full Name	tPA, partial
NCBI Official Synonym Full Names	plasminogen activator, tissue type
NCBI Official Symbol	PLAT
NCBI Official Synonym Symbols	TPA; T-PA
NCBI Protein Information	tissue-type plasminogen activator
UniProt Protein Name	TPA protein
UniProt Gene Name	tPA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/tissue-polypeptide-antigen-tpa/183059>