

Mouse Complement Component 4a (C4a) ELISA Kit | C4a elisa kit

Mouse Complement Component 4a (C4a) ELISA Kit

CATALOG #
AAA183328

REACTIVITY
Mouse


PRODUCT SYNONYMS

Complement Component 4a (C4a); Mouse Complement Component 4a (C4a) ELISA Kit; C4a anaphylatoxin; C4a elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Complement Component 4a (C4a). No significant cross-reactivity or interference between Complement Component 4a (C4a) and analogues was observed.
Sequence Length	1744
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates and Other Biological Fluids
Detection Range	31.2-2,000pg/mL
Sensitivity	< 11.9pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Complement Component 4a (C4a) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Complement Component 4a (C4a) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for C4a elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Complement Component 4a (C4a). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Complement Component 4a (C4a). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Complement Component 4a (C4a), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Complement Component 4a (C4a) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	67190748
NCBI GeneID	720
NCBI Accession #	NP_009224.2
NCBI GenBank Nucleotide #	NM_007293.2
UniProt Accession #	P0C0L4; P01028; P78445; Q13160; Q13906; A6H8M8; A6NHJ5; A7E2V2; B0QZR6; B0V2C8; B2RUT6; B7ZVZ6
Molecular Weight	187,704 Da
NCBI Official Full Name	complement C4-A isoform 1 preproprotein
NCBI Official Synonym Full Names	complement C4A (Rodgers blood group)
NCBI Official Symbol	C4A
NCBI Official Synonym Symbols	C4; RG; C4S; CO4; C4A2; C4A3; C4A4; C4A6; C4AD; CPAMD2
NCBI Protein Information	complement C4-A
UniProt Protein Name	Complement C4-A
UniProt Gene Name	C4A
UniProt Synonym Gene Names	CO4; CPAMD2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.