

Bovine Osteocalcin (OC) ELISA Kit | OC elisa kit

Cattle Osteocalcin (OC) ELISA Kit

CATALOG #
AAA183417

REACTIVITY
Bovine


PRODUCT SYNONYMS

Osteocalcin (OC); Cattle Osteocalcin (OC) ELISA Kit; BGLAP; OT; BGP; Bone Gla Protein; Bone Gamma-Carboxyglutamate Protein; Gamma-carboxyglutamic acid-containing protein; OC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Bovine
Specificity	This assay has high sensitivity and excellent specificity for detection of Osteocalcin (OC). No significant cross-reactivity or interference between Osteocalcin (OC) and analogues was observed.
Sequence Length	100
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates and Other Biological Fluids
Detection Range	78-5,000pg/mL
Sensitivity	< 32pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Osteocalcin (OC) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Osteocalcin (OC) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for OC elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Osteocalcin (OC). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Osteocalcin (OC). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Osteocalcin (OC), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Osteocalcin (OC) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	40316933
NCBI GeneID	632
NCBI Accession #	NP_954642.1
NCBI GenBank Nucleotide #	NM_199173.5
UniProt Accession #	P02818; Q5TCK6
Molecular Weight	10,963 Da
NCBI Official Full Name	osteocalcin preproprotein
NCBI Official Synonym Full Names	bone gamma-carboxyglutamate protein
NCBI Official Symbol	BGLAP
NCBI Official Synonym Symbols	OC; BGP; OCN
NCBI Protein Information	osteocalcin
UniProt Protein Name	Osteocalcin
UniProt Gene Name	BGLAP
UniProt Synonym Gene Names	BGP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.