

Rat Tissue Factor (TF) ELISA Kit | TF elisa kit

Rat Tissue Factor (TF) ELISA Kit

CATALOG #
AAA183476

REACTIVITY
Rat


PRODUCT SYNONYMS

Tissue Factor (TF); Rat Tissue Factor (TF) ELISA Kit; CD142; FIII; F3; TFA; Thromboplastin; Coagulation Factor III; TF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Tissue Factor (TF). No significant cross-reactivity or interference between Tissue Factor (TF) and analogues was observed.
Sequence Length	295
Assay Type	Double-antibody Sandwich
Samples	Plasma, Tissue Homogenates, Cell Lysates and Other Biological Fluids
Detection Range	31.2-2000pg/mL
Sensitivity	< 12.4pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Tissue Factor (TF) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Tissue Factor (TF) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TF elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Tissue Factor (TF). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Tissue Factor (TF). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Tissue Factor (TF), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Tissue Factor (TF) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	339508
NCBI GeneID	2152
UniProt Accession #	P13726; Q6FHG2; Q86WH4; D3DT47
Molecular Weight	27,145 Da
NCBI Official Full Name	tissue factor
NCBI Official Synonym Full Names	coagulation factor III, tissue factor
NCBI Official Symbol	F3
NCBI Official Synonym Symbols	TF; TFA; CD142
NCBI Protein Information	tissue factor
UniProt Protein Name	Tissue factor
UniProt Gene Name	F3
UniProt Synonym Gene Names	TF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/tissue-factor-tf/183476>