

Human Apolipoprotein E (APOE) ELISA Kit | APOE elisa kit

Human Apolipoprotein E (APOE) ELISA Kit

CATALOG #
AAA183838

REACTIVITY
Human


PRODUCT SYNONYMS

Apolipoprotein E (APOE); Human Apolipoprotein E (APOE) ELISA Kit; Apo-E; AD2; Apoprotein; Alzheimer Disease 2(E4-Associated; Late Onset; APOE elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Apolipoprotein E (APOE). No significant cross-reactivity or interference between Apolipoprotein E (APOE) and analogues was observed.
Sequence Length	317
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates and Other Biological Fluids
Detection Range	4.7-300ng/mL
Sensitivity	< 1.9ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Apolipoprotein E (APOE) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Apolipoprotein E (APOE) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for APOE elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Apolipoprotein E (APOE). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Apolipoprotein E (APOE). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Apolipoprotein E (APOE), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Apolipoprotein E (APOE) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557325
NCBI GeneID	348
NCBI Accession #	NP_000032.1
NCBI GenBank Nucleotide #	NM_000041.3
UniProt Accession #	P02649; Q9P2S4; B2RC15; C0JYY5
Molecular Weight	36,154 Da
NCBI Official Full Name	apolipoprotein E isoform b
NCBI Official Synonym Full Names	apolipoprotein E
NCBI Official Symbol	APOE
NCBI Official Synonym Symbols	AD2; LPG; APO-E; ApoE4; LDLCQ5
NCBI Protein Information	apolipoprotein E
UniProt Protein Name	Apolipoprotein E
UniProt Gene Name	APOE
UniProt Synonym Gene Names	Apo-E

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.