

Human Lymphotactin (Lptn) ELISA Kit | Lptn elisa kit

Human Lymphotactin (Lptn) ELISA Kit

CATALOG #
AAA183980

REACTIVITY
Human


PRODUCT SYNONYMS

Lymphotactin (Lptn); Human Lymphotactin (Lptn) ELISA Kit; XCL1; SCYC1; LTN; ATAC; SCM1-A; SCM1a; LPTN; Lymphotaxin; Chemokine C-Motif Ligand 1; Small Inducible Cytokine Subfamily C; Member 1; XC chemokine ligand 1; Lptn elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Lymphotactin (Lptn). No significant cross-reactivity or interference between Lymphotactin (Lptn) and analogues was observed.
Sequence Length	114
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates and Other Biological Fluids
Detection Range	15.6-1000pg/mL
Sensitivity	< 5.6pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Lymphotactin (Lptn) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Lymphotactin (Lptn) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Lptn elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Lymphotactin (Lptn). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Lymphotactin (Lptn). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Lymphotactin (Lptn), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Lymphotactin (Lptn) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506853
NCBI GeneID	6375
NCBI Accession #	NP_002986.1
NCBI GenBank Nucleotide #	NM_002995.2
UniProt Accession #	P47992; Q52MA8
Molecular Weight	12,517 Da
NCBI Official Full Name	lymphotactin
NCBI Official Synonym Full Names	X-C motif chemokine ligand 1
NCBI Official Symbol	XCL1
NCBI Official Synonym Symbols	LTN; ATAC; LPTN; SCM1; SCM-1; SCM1A; SCYC1; SCM-1a
NCBI Protein Information	lymphotactin
UniProt Protein Name	Lymphotactin
UniProt Gene Name	XCL1
UniProt Synonym Gene Names	LTN; SCYC1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/lymphotactin-lptn/183980>