

Mouse Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14) ELISA Kit | TNFSF14 elisa kit

Mouse Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14) ELISA Kit

CATALOG # AAA184035	REACTIVITY Mouse
-------------------------------	----------------------------


PRODUCT SYNONYMS

Tumor Necrosis Factor Ligand Superfamily; Member 14 (TNFSF14); Mouse Tumor Necrosis Factor Ligand Superfamily; Member 14 (TNFSF14) ELISA Kit; CD258; TR2; HVEM; LIGHT; LTg; Herpes virus entry mediator ligand; Herpesvirus entry mediator ligand; TNFSF14 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14). No significant cross-reactivity or interference between Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14) and analogues was observed.
Sequence Length	240
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Saliva, Tissue Homogenates and Other Biological Fluids
Detection Range	15.6-1000pg/mL
Sensitivity	< 6.0pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TNFSF14 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	25952144
NCBI GeneID	8740
NCBI Accession #	NP_003798.2
NCBI GenBank Nucleotide #	NM_003807.4
UniProt Accession #	O43557; O75476; Q6FHA1; Q8WVF8; Q96LD2; A8K7M2; C9J5H4
Molecular Weight	22,395 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 14 isoform 1
NCBI Official Synonym Full Names	TNF superfamily member 14
NCBI Official Symbol	TNFSF14
NCBI Official Synonym Symbols	LTg; CD258; HVEM-L; LIGHT
NCBI Protein Information	tumor necrosis factor ligand superfamily member 14
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 14
UniProt Gene Name	TNFSF14
UniProt Synonym Gene Names	HVEM-L; LIGHT; HVEM-L; Herpesvirus entry mediator ligand

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.