

Human Cluster Of Differentiation 160 (CD160) ELISA Kit | CD160 elisa kit

Human Cluster Of Differentiation 160 (CD160) ELISA Kit

CATALOG #
AAA184240

REACTIVITY
Human


PRODUCT SYNONYMS

Cluster Of Differentiation 160 (CD160); Human Cluster Of Differentiation 160 (CD160) ELISA Kit; BY55; NK1; NK28; CD160 Antigen; Natural killer cell receptor BY55; CD160 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of CD160. No significant cross-reactivity or interference between CD160 and analogues was observed.
Sequence Length	181
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	78-5,000pg/mL
Sensitivity	< 33pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CD160 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CD160 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD160 elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to CD160. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to CD160. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CD160, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of CD160 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CD160 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	5901910
NCBI GeneID	11126
NCBI Accession #	NP_008984.1
NCBI GenBank Nucleotide #	NM_007053.3
UniProt Accession #	O95971
Molecular Weight	19,810 Da
NCBI Official Full Name	CD160 antigen
NCBI Official Synonym Full Names	CD160 molecule
NCBI Official Symbol	CD160
NCBI Official Synonym Symbols	NK1; BY55; NK28
NCBI Protein Information	CD160 antigen
UniProt Protein Name	CD160 antigen
UniProt Gene Name	CD160
UniProt Synonym Gene Names	BY55

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/cluster-of-differentiation-160-cd160/184240>