

Human Fibrinogen Alpha (FGa) ELISA Kit | FGa elisa kit

Human Fibrinogen Alpha (FGa) ELISA Kit

CATALOG #
AAA184396

REACTIVITY
Human


PRODUCT SYNONYMS

Fibrinogen Alpha (FGa); Human Fibrinogen Alpha (FGa) ELISA Kit; Fib2; FG-A; Fibrinopeptide A; FGa elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Fibrinogen Alpha (FGa). No significant cross-reactivity or interference between Fibrinogen Alpha (FGa) and analogues was observed.
Sequence Length	644
Assay Type	Double-antibody Sandwich
Samples	Plasma
Detection Range	15.6-1,000ng/mL
Sensitivity	< 6.5ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Fibrinogen Alpha (FGa) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Fibrinogen Alpha (FGa) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FGa elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Fibrinogen Alpha (FGa). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Fibrinogen Alpha (FGa). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Fibrinogen Alpha (FGa), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Fibrinogen Alpha (FGa) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	11761629
NCBI GeneID	2243
NCBI Accession #	NP_068657.1
NCBI GenBank Nucleotide #	NM_021871.3
UniProt Accession #	P02671; Q4QQH7; Q9BX62; Q9UCH2; A8K3E4; D3DP14; D3DP15
Molecular Weight	69,757 Da
NCBI Official Full Name	fibrinogen alpha chain isoform alpha
NCBI Official Synonym Full Names	fibrinogen alpha chain
NCBI Official Symbol	FGA
NCBI Official Synonym Symbols	Fib2
NCBI Protein Information	fibrinogen alpha chain
UniProt Protein Name	Fibrinogen alpha chain
UniProt Gene Name	FGA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.