

Rat Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5) ELISA Kit | TNFRSF5 elisa kit

Rat Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5) ELISA Kit

CATALOG # AAA184496	REACTIVITY Rat
-------------------------------	--------------------------


PRODUCT SYNONYMS

Tumor Necrosis Factor Receptor Superfamily; Member 5 (TNFRSF5); Rat Tumor Necrosis Factor Receptor Superfamily; Member 5 (TNFRSF5) ELISA Kit; CD40; CDW40; Bp50; P50; B-cell surface antigen CD40; CD40L receptor; TNFRSF5 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5). No significant cross-reactivity or interference between Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5) and analogues was observed.
Sequence Length	203
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma and Other Biological Fluids
Detection Range	15.62-1000pg/mL
Sensitivity	< 5.2pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TNFRSF5 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Tumor Necrosis Factor Receptor Superfamily, Member 5 (TNFRSF5) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	23312371
NCBI GeneID	958
NCBI Accession #	NP_690593.1
NCBI GenBank Nucleotide #	NM_152854.3
UniProt Accession #	P25942; Q53GN5; Q5JY15; Q5U007; Q7M4Q8; Q86YK5; Q9BYU0; E1P5S9
Molecular Weight	22,259 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 5 isoform 2
NCBI Official Synonym Full Names	CD40 molecule
NCBI Official Symbol	CD40
NCBI Official Synonym Symbols	p50; Bp50; CDW40; TNFRSF5
NCBI Protein Information	tumor necrosis factor receptor superfamily member 5
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 5
UniProt Gene Name	CD40
UniProt Synonym Gene Names	TNFRSF5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.