

Human Complement C4-B (C4B) ELISA Kit | C4B elisa kit

Human Complement C4-B (C4B) ELISA Kit

CATALOG #
AAA184560

REACTIVITY
Human


PRODUCT SYNONYMS

Complement C4-B (C4B); Human Complement C4-B (C4B) ELISA Kit; CO4; CPAMD3; Basic complement C4; C3 and PZP-like alpha-2-macroglobulin domain-containing protein 3; C4B elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Complement C4-B (C4B). No significant cross-reactivity or interference between Complement C4-B (C4B) and analogues was observed.
Sequence Length	1744
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma and Other Biological Fluids
Detection Range	4.7-300ng/mL
Sensitivity	< 2.1ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Complement C4-B (C4B) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Complement C4-B (C4B) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for C4B elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Complement C4-B (C4B). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Complement C4-B (C4B). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Complement C4-B (C4B), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Complement C4-B (C4B) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	178557739
NCBI GeneID	721
NCBI Accession #	NP_001002029.3
NCBI GenBank Nucleotide #	NM_001002029.3
UniProt Accession #	P0C0L5; P01028; P78445; Q13160; Q13906; Q14033; Q14835; Q6U2E9; Q6U2G1; Q6U2I5; Q6U2L1; A2BHY4
Molecular Weight	192,751 Da
NCBI Official Full Name	complement C4-B preproprotein
NCBI Official Synonym Full Names	complement C4B (Chido blood group)
NCBI Official Symbol	C4B
NCBI Official Synonym Symbols	CH; C4F; CO4; C4B1; C4B2; C4B3; C4B5; C4BD; C4B12; C4B_2; CPAMD3
NCBI Protein Information	complement C4-B
UniProt Protein Name	Complement C4-B
UniProt Gene Name	C4B
UniProt Synonym Gene Names	CO4; CPAMD3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.