

Steroid 21-hydroxylase (CYP21A2) Recombinant Protein | CYP21A2 recombinant protein

Recombinant Human Steroid 21-hydroxylase (CYP21A2)

CATALOG # AAA18457	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
------------------------------	---


PRODUCT SYNONYMS

Steroid 21-hydroxylase (CYP21A2); Recombinant Human Steroid 21-hydroxylase (CYP21A2); 21-OHase; Cytochrome P-450c21; Cytochrome P450 21; Cytochrome P450 XXI; Cytochrome P450-C21; Cytochrome P450-C21B; CYP21A2 recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	1-494a; Full Length
Sequence	MLLLGLLLLP LLAGARLLWN WWKLRS LHLP PLAPGFLHLL QPDLPIYLLG LTQKFGPIYR LHLGLQDVVV LNSKRTIEEA MVKKWADFAG RPEPLTYKLV SKNYPDL SLG DYSLLWKAHK KLTRSALLG IRDSMEPVVE QLTQEFCERM RAQPGTPVAI EEEFSLTCS IICYLTFGDK IKDDNLMPAY YKCIQEV LKT WSHWSIQIVD VIPFLRFFPN PGLRRLKQAI EKR DHIVEMQ LRQH KESLVA GQWRDMMDYM LQGVAQPSME EGSGQLLEGH VHMAAVDLLI GGTETTANTL SWAVVFLHH PEIQRLQEE LDHELGP GAS SSRVPYKDRA RLPLLNATIA EVLRLRPVVP LALPHRTTRP SSISGYDIPE GTVIIPNLQG AHLDET VWER PHEFWPDRFL EPGKNSRALA FGCGARVCLG EPLARLELFV VLTRLLQAFT LLPSGDALPS LQPLPHCSVI LKMQPFQVRL QPRGMGAHSP GQNQ
Preparation and Storage	Store at -20 degree C, for extended storage, conserve at -20 degree C or -80 degree C.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for CYP21A2 recombinant protein	Specifically catalyzes the 21-hydroxylation of steroids. Required for the adrenal synthesis of mineralocorticoids and glucocorticoids.
Product Categories/Family for CYP21A2 recombinant protein	Cancer

References

Complete nucleotide sequence of two steroid 21-hydroxylase genes tandemly arranged in human chromosome a pseudogene and a genuine gene.Higashi Y., Yoshioka H., Yamane M., Gotoh O., Fujii-Kuriyama Y.Proc. Natl. Acad. Sci. U.S.A. 83:2841-2845(1986) Structure of human steroid 21-hydroxylase genes.White P.C., New M.I., Dupont B.Proc. Natl. Acad. Sci. U.S.A. 83:5111-5115(1986) Molecular characterization of the HLA-linked steroid 21-hydroxylase B gene from an individual with congenital adrenal hyperplasia.Rodrigues N.R., Dunham I., Yu C.Y., Carroll M.C., Porter R.R., Campbell R.D.EMBO J. 6:1653-1661(1987) Nonsense mutation causing steroid 21-hydroxylase deficiency.Globerman H., Amor M., Parker K.L., New M.I., White P.C.J. Clin. Invest. 82:139-144(1988) R339H and P453S CYP21 mutations associated with nonclassic steroid 21-hydroxylase deficiency that are not apparent gene conversions.Helmberg A., Tusie-Luna M.-T., Tabarelli M., Kofler R., White P.C.Mol. Endocrinol. 6:1318-1322(1992) Linkage analysis of the C4A/C4B copy number variation and polymorphisms of the adjacent steroid 21-hydroxylase gene in a healthy population.Blasko B., Banlaki Z., Gyapay G., Pozsonyi E., Sasvari-Szekely M., Rajczy K., Fust G., Szilagyi A.Mol. Immunol. 46:2623-2629(2009) Complete sequencing and characterization of 21,243 full-length human cDNAs.Ota T., Suzuki Y., Nishikawa T., Otsuki T., Sugiyama T., Irie R., Wakamatsu A., Hayashi K., Sato H., Nagai K., Kimura K., Makita H., Sekine M., Obayashi M., Nishi T., Shibahara T., Tanaka T., Ishii S., Yamamoto J., Saito K., Kawai Y., Isono Y., Nakamura Y., Nagahari K., Murakami K., Yasuda T., Iwayanagi T., Wagatsuma M., Shiratori A., Sudo H., Hosoiri T., Kaku Y., Kodaira H., Kondo H., Sugawara M., Takahashi M., Kanda K., Yokoi T., Furuya T., Kikkawa E., Omura Y., Abe K., Kamihara K., Katsuta N., Sato K., Tanikawa M., Yamazaki M., Ninomiya K., Ishibashi T., Yamashita H., Murakawa K., Fujimori K., Tanai H., Kimata M., Watanabe M., Hiraoka S., Chiba Y., Ishida S., Ono Y., Takiguchi S., Watanabe S., Yosida M., Hotuta T., Kusano J., Kanehori K., Takahashi-Fujii A., Hara H., Tanase T.-O., Nomura Y., Togiya S., Komai F., Hara R., Takeuchi K., Arita M., Imose N., Musashino K., Yuuki H., Oshima A., Sasaki N., Aotsuka S., Yoshikawa Y., Matsunawa H., Ichihara T., Shiohata N., Sano S., Moriya S., Momiyama H., Satoh N., Takami S., Terashima Y., Suzuki O., Nakagawa S., Senoh A., Mizoguchi H., Goto Y., Shimizu F., Wakebe H., Hishigaki H., Watanabe T., Sugiyama A., Takemoto M., Kawakami B., Yamazaki M., Watanabe K., Kumagai A., Itakura S., Fukuzumi Y., Fujimori Y., Komiyama M., Tashiro H., Tanigami A., Fujiwara T., Ono T., Yamada K., Fujii Y., Ozaki K., Hirao M., Ohmori Y., Kawabata A., Hikiji T., Kobatake N., Inagaki H., Ikema Y., Okamoto S., Okitani R., Kawakami T., Noguchi S., Itoh T., Shigeta K., Senba T., Matsumura K., Nakajima Y., Mizuno T., Morinaga M., Sasaki M., Togashi T., Oyama M., Hata H., Watanabe M., Komatsu T., Mizushima-Sugano J., Satoh T., Shirai Y., Takahashi Y., Nakagawa K., Okumura K., Nagase T., Nomura N., Kikuchi H., Masuho Y., Yamashita R., Nakai K., Yada T., Nakamura Y., Ohara O., Isogai T., Sugano S.Nat. Genet. 36:40-45(2004) The DNA sequence and analysis of human chromosome 6.Mungall A.J., Palmer S.A., Sims S.K., Edwards C.A., Ashurst J.L., Wilming L., Jones M.C., Horton R., Hunt S.E., Scott C.E., Gilbert J.G.R., Clamp M.E., Bethel G., Milne S., Ainscough R., Almeida J.P., Ambrose K.D., Andrews T.D., Ashwell R.I.S., Babbage A.K., Bagguley C.L., Bailey J., Banerjee R., Barker D.J., Barlow K.F., Bates K., Beare D.M., Beasley H., Beasley O., Bird C.P., Blakey S.E., Bray-Allen S., Brook J., Brown A.J., Brown J.Y., Burford D.C., Burrill W., Burton J., Carder C., Carter N.P., Chapman J.C., Clark S.Y., Clark G., Clee C.M., Clegg S., Cobley V., Collier R.E., Collins J.E., Colman L.K., Corby N.R., Coville G.J., Culley K.M., Dhani P., Davies J., Dunn M., Earthrowl M.E., Ellington A.E., Evans K.A., Faulkner L., Francis M.D., Frankish A., Frankland J., French L., Garner P., Garnett J., Ghorri M.J., Gilby L.M., Gillson C.J., Glithero R.J., Grafham D.V., Grant M., Gribble S., Griffiths C., Griffiths M.N.D., Hall R., Halls K.S., Hammond S., Harley J.L., Hart E.A., Heath P.D., Heathcott R., Holmes S.J., Howden P.J., Howe K.L., Howell G.R., Huckle E., Humphray S.J., Humphries M.D., Hunt A.R., Johnson C.M., Joy A.A., Kay M., Keenan S.J., Kimberley A.M., King A., Laird G.K., Langford C., Lawlor S., Leongamornlert D.A., Leversha M., Lloyd C.R., Lloyd D.M., Loveland J.E., Lovell J., Martin S., Mashreghi-Mohammadi M., Maslen G.L., Matthews L., McCann O.T., McLaren S.J., McLay K., McMurray A., Moore M.J.F., Mullikin J.C., Niblett D., Nickerson T., Novik K.L., Oliver K., Overton-Larty E.K., Parker A., Patel R., Pearce A.V., Peck A.I., Phillimore B.J.C.T., Phillips S., Plumb R.W., Porter K.M., Ramsey Y., Ranby S.A., Rice C.M., Ross M.T., Searle S.M., Sehra H.K., Sheridan E., Skuce C.D., Smith S., Smith M., Spraggon L., Squares S.L., Steward C.A., Sycamore N., Tamlyn-Hall G., Tester J., Theaker A.J., Thomas D.W., Thorpe A., Tracey A., Tromans A., Tubby B., Wall M., Wallis J.M., West A.P., White S.S., Whitehead S.L., Whittaker H., Wild A., Willey D.J., Wilmer T.E., Wood J.M., Wray P.W., Wyatt J.C., Young L., Younger R.M., Bentley D.R., Coulson A., Durbin R.M., Hubbard T., Sulston J.E., Dunham I., Rogers J., Beck S.Nature 425:805-811(2003) Mural R.J., Istrail S., Sutton G., Florea L., Halpern A.L., Mobarry C.M., Lippert R., Walenz B., Shatkay H., Dew I., Miller J.R., Flanigan M.J., Edwards N.J., Bolanos R., Fasulo D., Halldorsson B.V., Hannonhalli S., Turner R., Yooshep S., Lu F., Nusskern D.R., Shue B.C., Zheng X.H., Zhong F., Delcher A.L., Huson D.H., Kravitz S.A., Mouchard L., Reinert K., Remington K.A., Clark A.G., Waterman M.S., Eichler E.E., Adams M.D., Hunkapiller M.W., Myers E.W., Venter J.C. A de novo pathological point mutation at the 21-hydroxylase locus implications for gene conversion in the human genome.Collier S., Tassabehji M., Sinnott P., Strachan T.Nat. Genet. 3:260-265(1993) Mapping of steroid 21-hydroxylase genes adjacent to complement component C4 genes in HLA, the major histocompatibility complex in man.Carroll M.C., Campbell R.D., Porter R.R.Proc. Natl. Acad. Sci. U.S.A. 82:521-525(1985) Mutation in the CYP21B gene (Ile-172-->Asn) causes steroid 21-hydroxylase deficiency.Amor M., Parker K.L., Globerman H., New M.I., White P.C.Proc. Natl. Acad. Sci. U.S.A. 85:1600-1604(1988) P450XXI (steroid 21-hydroxylase) gene deletions are not found in family studies of congenital adrenal hyperplasia.Matteson K.J., Phillips J.A. III, Miller W.L., Chung B.C., Orlando P.J., Frisch H., Ferrandez A., Burr I.M.Proc. Natl. Acad. Sci. U.S.A. 84:5858-5862(1987) Mutations of P450c21 (steroid 21-hydroxylase) at Cys428, Val281, and Ser268 result in complete, partial, or no loss of enzymatic activity, respectively.Wu D.-A., Chung B.-C.J. Clin. Invest. 88:519-523(1991) Molecular genetics of 21-hydroxylase deficient late-onset adrenal hyperplasia.Gunn S.K., Sherman L.D., Therrell B.L., Owerbach D.I.Semin. Reprod. Endocrinol. 11:347-352(1993) Mutations in steroid 21-hydroxylase (CYP21).White P.C., Tusie-Luna M.-T., New M.I., Speiser P.W.Hum. Mutat. 3:373-378(1994) Molecular genetic analysis of nonclassic steroid 21-hydroxylase deficiency associated with HLA-B14,DR1.Speiser P.W., New M.I., White P.C.N. Engl. J. Med. 319:19-23(1988) A missense mutation at Ile172-->Asn or Arg356-->Trp causes steroid 21-hydroxylase deficiency.Chiou S.-H., Hu M.-C., Chung B.-C.J. Biol. Chem. 265:3549-3552(1990) Substitution of

NCBI AND UNIPROT INFORMATION

NCBI GI #	323510663
NCBI GeneID	1589
NCBI Accession #	NP_000491.4
NCBI GenBank Nucleotide #	NM_000500.7
UniProt Accession #	P08686; P04033; Q01204; Q08AG8; Q16749; Q16806; Q5ST44; Q96NU8; A2BHY6
Molecular Weight	57.9 kDa
NCBI Official Full Name	steroid 21-hydroxylase isoform a
NCBI Official Synonym Full Names	cytochrome P450 family 21 subfamily A member 2
NCBI Official Symbol	CYP21A2
NCBI Official Synonym Symbols	CAH1; CPS1; CA21H; CYP21; CYP21B; P450c21B
NCBI Protein Information	steroid 21-hydroxylase
UniProt Protein Name	Steroid 21-hydroxylase
UniProt Gene Name	CYP21A2
UniProt Synonym Gene Names	CYP21; CYP21B
UniProt Entry Name	CP21A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/cyp21a2-recombinant-protein-e-coli-or-yeast-or-baculovirus-or-mammalian-cell/18457>