

Human Lymphocyte Activation Gene 3 (LAG3) ELISA Kit | LAG3 elisa kit

Human Lymphocyte Activation Gene 3 (LAG3) ELISA Kit

CATALOG #
AAA184935

REACTIVITY
Human


PRODUCT SYNONYMS

Lymphocyte Activation Gene 3 (LAG3); Human Lymphocyte Activation Gene 3 (LAG3) ELISA Kit; CD223; Protein FDC; LAG3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Lymphocyte Activation Gene 3 (LAG3). No significant cross-reactivity or interference between Lymphocyte Activation Gene 3 (LAG3) and analogues was observed.
Sequence Length	360
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates, Cell Lysates and Other Biological Fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.056ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Lymphocyte Activation Gene 3 (LAG3) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Lymphocyte Activation Gene 3 (LAG3) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for LAG3 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Lymphocyte Activation Gene 3 (LAG3). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Lymphocyte Activation Gene 3 (LAG3). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Lymphocyte Activation Gene 3 (LAG3), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Lymphocyte Activation Gene 3 (LAG3) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	167614500
NCBI GeneID	3902
NCBI Accession #	NP_002277.4
NCBI GenBank Nucleotide #	NM_002286.5
UniProt Accession #	P18627; Q7Z643; A8K7T9
Molecular Weight	39,103 Da
NCBI Official Full Name	lymphocyte activation gene 3 protein
NCBI Official Synonym Full Names	lymphocyte activating 3
NCBI Official Symbol	LAG3
NCBI Official Synonym Symbols	CD223
NCBI Protein Information	lymphocyte activation gene 3 protein
UniProt Protein Name	Lymphocyte activation gene 3 protein
UniProt Gene Name	LAG3
UniProt Synonym Gene Names	FDC; LAG-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.