

Mouse B-Cell Activating Factor (BAFF) ELISA Kit | BAFF elisa kit

Mouse B-Cell Activating Factor (BAFF) ELISA Kit

CATALOG #
AAA184960

REACTIVITY
Mouse


PRODUCT SYNONYMS

B-Cell Activating Factor (BAFF); Mouse B-Cell Activating Factor (BAFF) ELISA Kit; CD257; TNFSF13B; BLYS; TALL1; THANK; TNFSF20; ZTNF4; B lymphocyte stimulator; Tumor Necrosis Factor Ligand Superfamily Member 13B; Dendritic cell-derived TNF-like; BAFF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of B-Cell Activating Factor (BAFF). No significant cross-reactivity or interference between B-Cell Activating Factor (BAFF) and analogues was observed.
Sequence Length	164
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates and Other Biological Fluids
Detection Range	12.5-800pg/mL
Sensitivity	< 4.9pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level B-Cell Activating Factor (BAFF) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level B-Cell Activating Factor (BAFF) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for BAFF elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to B-Cell Activating Factor (BAFF). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to B-Cell Activating Factor (BAFF). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain B-Cell Activating Factor (BAFF), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of B-Cell Activating Factor (BAFF) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4585275
NCBI GeneID	10673
UniProt Accession #	Q9Y275; Q6FHD6; Q7Z5J2; E0ADT7
Molecular Weight	17,578 Da
NCBI Official Full Name	B-cell activating factor
NCBI Official Synonym Full Names	TNF superfamily member 13b
NCBI Official Symbol	TNFSF13B
NCBI Official Synonym Symbols	DTL; BAFF; BLYS; CD257; TALL1; THANK; ZTNF4; TALL-1; TNLG7A; TNFSF20
NCBI Protein Information	tumor necrosis factor ligand superfamily member 13B
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 13B
UniProt Gene Name	TNFSF13B
UniProt Synonym Gene Names	BAFF; BLYS; TALL1; TNFSF20; ZTNF4; BLyS; TALL-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.