

Human Mucin 7, Secreted (MUC7) ELISA Kit | MUC7 elisa kit

Human Mucin 7, Secreted (MUC7) ELISA Kit

CATALOG #
AAA185088

REACTIVITY
Human


PRODUCT SYNONYMS

Mucin 7; Secreted (MUC7); Human Mucin 7; Secreted (MUC7) ELISA Kit; MG2; Salivary; Apo-MG2; Salivary mucin-7; MUC7 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of MUC7. No significant cross-reactivity or interference between MUC7 and analogues was observed.
Sequence Length	377
Assay Type	Quantitative Sandwich
Samples	Human saliva, tissue homogenates, cell lysates and other biological fluids
Detection Range	0.312-20ng/mL
Sensitivity	< 0.112ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level MUC7 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level MUC7 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MUC7 elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to MUC7. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to MUC7. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MUC7, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of MUC7 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of MUC7 in human saliva, tissue homogenates, cell lysates and other biological fluids.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	119626026
NCBI GeneID	4589
UniProt Accession #	Q8TAX7; Q9UCD7; Q9UCD8
Molecular Weight	39,159 Da
NCBI Official Full Name	mucin 7, secreted
NCBI Official Synonym Full Names	mucin 7, secreted
NCBI Official Symbol	MUC7
NCBI Official Synonym Symbols	MG2
NCBI Protein Information	mucin-7
UniProt Protein Name	Mucin-7
UniProt Gene Name	MUC7
UniProt Synonym Gene Names	MG2; MUC-7

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.