

Rat Cyclin Dependent Kinase 2 (CDK2) ELISA Kit | CDK2 elisa kit

Rat Cyclin Dependent Kinase 2 (CDK2) ELISA Kit

CATALOG #
AAA185211

REACTIVITY
Rat


PRODUCT SYNONYMS

Cyclin Dependent Kinase 2 (CDK2); Rat Cyclin Dependent Kinase 2 (CDK2) ELISA Kit; CDKN2; p33(CDK2); Cell division protein kinase 2; p33 protein kinase; CDK2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Cyclin Dependent Kinase 2 (CDK2). No significant cross-reactivity or interference between Cyclin Dependent Kinase 2 (CDK2) and analogues was observed.
Sequence Length	â€
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates and Other Biological Fluids
Detection Range	0.312-20ng/mL
Sensitivity	< 0.133ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Cyclin Dependent Kinase 2 (CDK2) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Cyclin Dependent Kinase 2 (CDK2) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CDK2 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Cyclin Dependent Kinase 2 (CDK2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Cyclin Dependent Kinase 2 (CDK2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Cyclin Dependent Kinase 2 (CDK2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Cyclin Dependent Kinase 2 (CDK2) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	41054836
NCBI GeneID	362817
NCBI Accession #	NP_955795.1
NCBI GenBank Nucleotide #	NM_199501.1
UniProt Accession #	Q63699; O09136
Molecular Weight	~ Da
NCBI Official Full Name	cyclin-dependent kinase 2
NCBI Official Synonym Full Names	cyclin dependent kinase 2
NCBI Official Symbol	Cdk2
NCBI Protein Information	cyclin-dependent kinase 2
UniProt Protein Name	Cyclin-dependent kinase 2
UniProt Gene Name	Cdk2
UniProt Synonym Gene Names	Cdkn2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.