

Cellular tumor antigen p53 (TP53) Recombinant Protein | TP53 recombinant protein

Recombinant Human Cellular tumor antigen p53 (TP53)

CATALOG # AAA18532	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
------------------------------	---


PRODUCT SYNONYMS

Cellular tumor antigen p53 (TP53); Recombinant Human Cellular tumor antigen p53 (TP53); Antigen NY-CO-13; Phosphoprotein p53; Tumor suppressor p53; TP53 recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	1-393aa; Full Length
Sequence	MEEPQSDPSV EPPLSQETFS DLWKLLEN VLSPLPSQAM DDLMLSPDDI EQWFTEDPGP DEAPRMPEAA PPVAPAPAAP TPAAPAPAPS WPLSSSVPSQ KTYQGSYGFR LGFLHSGTAK SVTCTYSPAL NKMFCQLAKT CPVQLWVDST PPPGTRVRAM AIYKQSQHMT EVVRRCPHHE RCSDSGLAP PQHLIRVEGN LRVEYLDDRN TFRHSVVVPY EPPEVGSDCT TIHNYMCNS SCMGGMNRRP ILTIITLED SGNLLGRNSF EVRVCACPGR DRRTEENLR KKGEPHHELP PGSTKRALPN NTSSSPQPK KPLDGEYFTL QIRGRERFEM FRELNEALEL KDAQAGKEPG GSRAHSSHLK SKKGQSTSRH KKLMEFKTEGP DSD
Preparation and Storage	Store at -20 degree C, for extended storage, conserve at -20 degree C or -80 degree C.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for TP53 recombinant protein	Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction ses to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and se to have to effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-ARNTL/BMAL1-mediated transcriptional activation of PER2.
Product Categories/Family for TP53 recombinant protein	Apoptosis

References

Human p53 cellular tumor antigen cDNA sequence and expression in COS cells.Zakut-Houri R., Bienz-Tadmor B., Givol D., Oren M.EMBO J. 4:1251-1255(1985) Characterization of the human p53 gene.Lamb P., Crawford L.Mol. Cell. Biol. 6:1379-1385(1986) Molecular cloning and in vitro expression of a cDNA clone for human cellular tumor antigen p53.Harlow E., Williamson N.M., Ralston R., Helfman D.M., Adams T.E.Mol. Cell. Biol. 5:1601-1610(1985) Molecular basis for heterogeneity of the human p53 protein.Harris N., Brill E., Shohat O., Prokocimer M., Wolf D., Arai N., Rotter V.Mol. Cell. Biol. 6:4650-4656(1986) A variation in the structure of the protein-coding region of the human p53 gene.Buchman V.L., Chumakov P.M., Ninkina N.N., Samarina O.P., Georgiev G.P.Gene 70:245-252(1988) p53 is frequently mutated in Burkitt's lymphoma cell lines.Farrell P.J., Allan G., Shanahan F., Vousden K.H., Crook T.EMBO J. 10:2879-2887(1991) Isolation of two cell lines from a human malignant glioma specimen differing in sensitivity to radiation and chemotherapeutic drugs.Allalunis-Turner M.J., Barron G.M., Day R.S. III, Dobler K.D., Mirzayans R.Radiat. Res. 134:349-354(1993) Hyaluronidase induction of a WW domain-containing oxidoreductase that enhances tumor necrosis factor cytotoxicity.Chang N.-S., Pratt N., Heath J., Schultz L., Sleva D., Carey G.B., Zevotek N.J. Biol. Chem. 276:3361-3370(2001) p53 isoforms can regulate p53 transcriptional activity.Bourdon J.C., Fernandes K., Murray-Zmijewski F., Liu G., Diot A., Xirodimas D.P., Saville M.K., Lane D.P.Genes Dev. 19:2122-2137(2005) Chumakov P.M., Almazov V.P., Jenkins J.R.P53 genomic sequence. Corrections and polymorphism.Rozemuller E.H., Tilanus M.G.J.NIEHS SNPs programHuman TP53 from the malignant glioma-derived cell lines M059J and M059K has a cancer-associated mutation in exon 8.Anderson C.W., Allalunis-Turner M.J.Radiat. Res. 154:473-476(2000) Identification of a tumor-rejection antigen recognized by HLA-B46 restricted CTL.Azuma K., Shichijo S., Itoh K.Complete sequencing and characterization of 21,243 full-length human cDNAs.Ota T., Suzuki Y., Nishikawa T., Otsuki T., Sugiyama T., Irie R., Wakamatsu A., Hayashi K., Sato H., Nagai K., Kimura K., Makita H., Sekine M., Obayashi M., Nishi T., Shibahara T., Tanaka T., Ishii S., Yamamoto J., Saito K., Kawai Y., Isono Y., Nakamura Y., Nagahari K., Murakami K., Yasuda T., Iwayanagi T., Wagatsuma M., Shiratori A., Sudo H., Hosoiri T., Kaku Y., Kodaira H., Kondo H., Sugawara M., Takahashi M., Kanda K., Yokoi T., Furuya T., Kikkawa E., Omura Y., Abe K., Kamihara K., Katsuta N., Sato K., Tanikawa M., Yamazaki M., Ninomiya K., Ishibashi T., Yamashita H., Murakawa K., Fujimori K., Tanai H., Kimata M., Watanabe M., Hiraoka S., Chiba Y., Ishida S., Ono Y., Takiguchi S., Watanabe S., Yosida M., Hotuta T., Kusano J., Kanehori K., Takahashi-Fujii A., Hara H., Tanase T.-O., Nomura Y., Togiya S., Komai F., Hara R., Takeuchi K., Arita M., Imose N., Musashino K., Yuuki H., Oshima A., Sasaki N., Aotsuka S., Yoshikawa Y., Matsunawa H., Ichihara T., Shiohata N., Sano S., Moriya S., Momiyama H., Satoh N., Takami S., Terashima Y., Suzuki O., Nakagawa S., Senoh A., Mizoguchi H., Goto Y., Shimizu F., Wakebe H., Hishigaki H., Watanabe T., Sugiyama A., Takemoto M., Kawakami B., Yamazaki M., Watanabe K., Kumagai A., Itakura S., Fukuzumi Y., Fujimori Y., Komiyama M., Tashiro H., Tanigami A., Fujiwara T., Ono T., Yamada K., Fujii Y., Ozaki K., Hirao M., Ohmori Y., Kawabata A., Hikiji T., Kobatake N., Inagaki H., Ikema Y., Okamoto S., Okitani R., Kawakami T., Noguchi S., Itoh T., Shigeta K., Senba T., Matsumura K., Nakajima Y., Mizuno T., Morinaga M., Sasaki M., Togashi T., Oyama M., Hata H., Watanabe M., Komatsu T., Mizushima-Sugano J., Satoh T., Shirai Y., Takahashi Y., Nakagawa K., Okumura K., Nagase T., Nomura N., Kikuchi H., Masuho Y., Yamashita R., Nakai K., Yada T., Nakamura Y., Ohara O., Isogai T., Sugano S.Nat. Genet. 36:40-45(2004) DNA sequence of human chromosome 17 and analysis of rearrangement in the human lineage.Zody M.C., Garber M., Adams D.J., Sharpe T., Harrow J., Lupski J.R., Nicholson C., Searle S.M., Wilming L., Young S.K., Abouelleil A., Allen N.R., Bi W., Bloom T., Borowsky M.L., Bugalter B.E., Butler J., Chang J.L., Chen C.-K., Cook A., Corum B., Cuomo C.A., de Jong P.J., DeCaprio D., Dewar K., FitzGerald M., Gilbert J., Gibson R., Gnerre S., Goldstein S., Grafham D.V., Grocock R., Hafez N., Hagopian D.S., Hart E., Norman C.H., Humphray S., Jaffe D.B., Jones M., Kamal M., Khodiyar V.K., LaButti K., Laird G., Lehoczy J., Liu X., Lokytsang T., Loveland J., Lui A., Macdonald P., Major J.E., Matthews L., Mauceli E., McCarroll S.A., Mihalev A.H., Mudge J., Nguyen C., Nicol R., O'Leary S.B., Osoegawa K., Schwartz D.C., Shaw-Smith C., Stankiewicz P., Steward C., Swarbreck D., Venkataraman V., Whittaker C.A., Yang X., Zimmer A.R., Bradley A., Hubbard T., Birren B.W., Rogers J., Lander E.S., Nusbaum C.Nature 440:1045-1049(2006)

NCBI AND UNIPROT INFORMATION

NCBI GI #	120407068
NCBI GeneID	7157
NCBI Accession #	NP_000537.3
NCBI GenBank Nucleotide #	NM_000546.5
UniProt Accession #	P04637; Q15086; Q15087; Q15088; Q16535; Q16807; Q16808; Q16809; Q16810; Q16811; Q16848; Q2XN98
Molecular Weight	45.7 kDa
NCBI Official Full Name	cellular tumor antigen p53 isoform a
NCBI Official Synonym Full Names	tumor protein p53

NCBI Official Symbol	TP53
NCBI Official Synonym Symbols	P53; BCC7; LFS1; TRP53
NCBI Protein Information	cellular tumor antigen p53
UniProt Protein Name	Cellular tumor antigen p53
UniProt Gene Name	TP53
UniProt Synonym Gene Names	P53
UniProt Entry Name	P53_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/tp53-recombinant-protein-e-coli-or-yeast-or-baculovirus-or-mammalian-cell/18532>