

Human Hepatocyte Growth Factor Receptor (HGFR) ELISA Kit | HGFR elisa kit

Human Hepatocyte Growth Factor Receptor (HGFR) ELISA Kit

CATALOG #
AAA185579

REACTIVITY
Human


PRODUCT SYNONYMS

Hepatocyte Growth Factor Receptor (HGFR); Human Hepatocyte Growth Factor Receptor (HGFR) ELISA Kit; CMET; RCCP2; C-Met; Met Proto-Oncogene; Mesenchymal-Epithelial Transition Factor; HGF/SF receptor; Proto-oncogene c-Met; Scatter factor receptor; Tyrosine-protein kinase Met; HGFR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Hepatocyte Growth Factor Receptor (HGFR). No significant cross-reactivity or interference between Hepatocyte Growth Factor Receptor (HGFR) and analogues was observed.
Sequence Length	764
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates and Other Biological Fluids
Detection Range	125-8000pg/mL
Sensitivity	< 49pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Hepatocyte Growth Factor Receptor (HGFR) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Hepatocyte Growth Factor Receptor (HGFR) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HGFR elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Hepatocyte Growth Factor Receptor (HGFR). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Hepatocyte Growth Factor Receptor (HGFR). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Hepatocyte Growth Factor Receptor (HGFR), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Hepatocyte Growth Factor Receptor (HGFR) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	42741655
NCBI GeneID	4233
NCBI Accession #	NP_000236.2
NCBI GenBank Nucleotide #	NM_000245.3
UniProt Accession #	P08581; O60366; Q12875; Q9UDX7; Q9UPL8; A1L467; B5A932; E7EQ94
Molecular Weight	85,745 Da
NCBI Official Full Name	hepatocyte growth factor receptor isoform b preproprotein
NCBI Official Synonym Full Names	MET proto-oncogene, receptor tyrosine kinase
NCBI Official Symbol	MET
NCBI Official Synonym Symbols	HGFR; AUTS9; RCCP2; c-Met; DFNB97
NCBI Protein Information	hepatocyte growth factor receptor
UniProt Protein Name	Hepatocyte growth factor receptor
UniProt Gene Name	MET
UniProt Synonym Gene Names	HGF receptor; SF receptor

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.