

Mouse Reelin (RL) ELISA Kit | RL elisa kit

Mouse Reelin (RL) ELISA Kit

CATALOG #
AAA185924

REACTIVITY
Mouse


PRODUCT SYNONYMS

Reelin (RL); Mouse Reelin (RL) ELISA Kit; RELN; RL elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Reelin (RL). No significant cross-reactivity or interference between Reelin (RL) and analogues was observed.
Sequence Length	3,427
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates, Cell Lysates and Other Biological Fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.055ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Reelin (RL) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Reelin (RL) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for RL elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Reelin (RL). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Reelin (RL). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Reelin (RL), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Reelin (RL) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	1743885
NCBI GeneID	5649
UniProt Accession #	P78509; Q86UJ0; Q86UJ8; Q8NDV0; Q9UDQ2; A4D0P9; A4D0Q0
Molecular Weight	384,093 Da
NCBI Official Full Name	reelin
NCBI Official Synonym Full Names	reelin
NCBI Official Symbol	RELN
NCBI Official Synonym Symbols	RL; ETL7; LIS2; PRO1598
NCBI Protein Information	reelin
UniProt Protein Name	Reelin
UniProt Gene Name	RELN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.