

Human Transgelin (TAGLN) ELISA Kit | TAGLN elisa kit

Human Transgelin (TAGLN) ELISA Kit

CATALOG #
AAA185930

REACTIVITY
Human


PRODUCT SYNONYMS

Transgelin (TAGLN); Human Transgelin (TAGLN) ELISA Kit; SMCC; SM22; TAGLN1; WS3-10; Transgelin Variant 2; 22 kDa actin-binding protein; Protein WS3-10; Smooth muscle protein 22-alpha; TAGLN elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Transgelin (TAGLN). No significant cross-reactivity or interference between Transgelin (TAGLN) and analogues was observed.
Sequence Length	201
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates and Other Biological Fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.065ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Transgelin (TAGLN) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Transgelin (TAGLN) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TAGLN elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Transgelin (TAGLN). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Transgelin (TAGLN). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Transgelin (TAGLN), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Transgelin (TAGLN) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	48255907
NCBI GeneID	6876
NCBI Accession #	NP_001001522.1
NCBI GenBank Nucleotide #	NM_001001522.2
UniProt Accession #	Q01995; O15542
Molecular Weight	22,611 Da
NCBI Official Full Name	transgelin
NCBI Official Synonym Full Names	transgelin
NCBI Official Symbol	TAGLN
NCBI Official Synonym Symbols	SM22; SMCC; TAGLN1; WS3-10; SM22-alpha
NCBI Protein Information	transgelin
UniProt Protein Name	Transgelin
UniProt Gene Name	TAGLN
UniProt Synonym Gene Names	SM22; WS3-10; SM22-alpha

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.