

Mouse Fibroblast Growth Factor 17 (FGF17) ELISA Kit | FGF17 elisa kit

Mouse Fibroblast Growth Factor 17 (FGF17) ELISA Kit

CATALOG #
AAA185998

REACTIVITY
Mouse


PRODUCT SYNONYMS

Fibroblast Growth Factor 17 (FGF17); Mouse Fibroblast Growth Factor 17 (FGF17) ELISA Kit; FGF17 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Fibroblast Growth Factor 17 (FGF17). No significant cross-reactivity or interference between Fibroblast Growth Factor 17 (FGF17) and analogues was observed.
Sequence Length	205
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates and Other Biological Fluids
Detection Range	15.6-1,000pg/mL
Sensitivity	< 6.0pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Fibroblast Growth Factor 17 (FGF17) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Fibroblast Growth Factor 17 (FGF17) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FGF17 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Fibroblast Growth Factor 17 (FGF17). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Fibroblast Growth Factor 17 (FGF17). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Fibroblast Growth Factor 17 (FGF17), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Fibroblast Growth Factor 17 (FGF17) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	109731710
NCBI GeneID	8822
Molecular Weight	23,669 Da
NCBI Official Full Name	Fibroblast growth factor 17
NCBI Official Synonym Full Names	fibroblast growth factor 17
NCBI Official Symbol	FGF17
NCBI Official Synonym Symbols	HH20; FGF-13; FGF-17
NCBI Protein Information	fibroblast growth factor 17
UniProt Protein Name	Fibroblast growth factor 17
UniProt Gene Name	FGF17
UniProt Synonym Gene Names	FGF-17

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/fibroblast-growth-factor-17-fgf17/185998>