

Human Interleukin 17D (IL17D) ELISA Kit | IL17D elisa kit

Human Interleukin 17D (IL17D) ELISA Kit

CATALOG #
AAA186233

REACTIVITY
Human


PRODUCT SYNONYMS

Interleukin 17D (IL17D); Human Interleukin 17D (IL17D) ELISA Kit; Interleukin-27; IL17D elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Interleukin 17D (IL17D). No significant cross-reactivity or interference between Interleukin 17D (IL17D) and analogues was observed.
Sequence Length	202
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates and Other Biological Fluids
Detection Range	15.6-1,000pg/mL
Sensitivity	< 6.2pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Interleukin 17D (IL17D) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Interleukin 17D (IL17D) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL17D elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Interleukin 17D (IL17D). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Interleukin 17D (IL17D). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Interleukin 17D (IL17D), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Interleukin 17D (IL17D) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	19923715
NCBI GeneID	53342
NCBI Accession #	NP_612141.1
NCBI GenBank Nucleotide #	NM_138284.1
UniProt Accession #	Q8TAD2; B1AM69
Molecular Weight	21,893 Da
NCBI Official Full Name	interleukin-17D
NCBI Official Synonym Full Names	interleukin 17D
NCBI Official Symbol	IL17D
NCBI Official Synonym Symbols	IL-17D
NCBI Protein Information	interleukin-17D
UniProt Protein Name	Interleukin-17D
UniProt Gene Name	IL17D
UniProt Synonym Gene Names	IL-27

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.