

Human Testis Anion Transporter 1 (TAT1) ELISA Kit | TAT1 elisa kit

Human Testis Anion Transporter 1 (TAT1) ELISA Kit

CATALOG #
AAA186590

REACTIVITY
Human


PRODUCT SYNONYMS

Testis Anion Transporter 1 (TAT1); Human Testis Anion Transporter 1 (TAT1) ELISA Kit; SLC26A8; Solute Carrier Family 26 Member 8; Anion exchange transporter; TAT1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Testis Anion Transporter 1 (TAT1). No significant cross-reactivity or interference between Testis Anion Transporter 1 (TAT1) and analogues was observed.
Sequence Length	768
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates and Other Biological Fluids
Detection Range	0.312-20ng/mL
Sensitivity	< 0.118ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Testis Anion Transporter 1 (TAT1) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Testis Anion Transporter 1 (TAT1) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TAT1 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Testis Anion Transporter 1 (TAT1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Testis Anion Transporter 1 (TAT1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Testis Anion Transporter 1 (TAT1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Testis Anion Transporter 1 (TAT1) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	16418457
NCBI GeneID	116369
NCBI Accession #	NP_443193.1
NCBI GenBank Nucleotide #	NM_052961.3
UniProt Accession #	Q96RN1; Q5JVR5; Q812C7; Q8TC65; Q96MA0; Q96PK8
Molecular Weight	85,680 Da
NCBI Official Full Name	testis anion transporter 1 isoform a
NCBI Official Synonym Full Names	solute carrier family 26 member 8
NCBI Official Symbol	SLC26A8
NCBI Official Synonym Symbols	TAT1; SPGF3
NCBI Protein Information	testis anion transporter 1
UniProt Protein Name	Testis anion transporter 1
UniProt Gene Name	SLC26A8

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.