

Human 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) ELISA Kit | HSD17b3 elisa kit

Human 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) ELISA Kit

CATALOG # AAA186752	REACTIVITY Human
-------------------------------	----------------------------


PRODUCT SYNONYMS

17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3); Human 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) ELISA Kit; SDR12C2; EDH17B3; Short Chain Dehydrogenase/Reductase Family 12C; Member 2; Testosterone 17-Beta-Dehydrogenase 3; Testicular 17-beta-hydroxysteroid dehydrogenase; HSD17b3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3). No significant cross-reactivity or interference between 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) and analogues was observed.
Sequence Length	305
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates and Other Biological Fluids
Detection Range	15.62-1000pg/mL
Sensitivity	< 6.4pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HSD17b3 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	1513310
NCBI GeneID	15487
UniProt Accession #	P70385; G3UWF9
Molecular Weight	34,330 Da
NCBI Official Full Name	17-beta-hydroxysteroid dehydrogenase type 3
NCBI Official Synonym Full Names	hydroxysteroid (17-beta) dehydrogenase 3
NCBI Official Symbol	Hsd17b3
NCBI Protein Information	testosterone 17-beta-dehydrogenase 3
UniProt Protein Name	Testosterone 17-beta-dehydrogenase 3
UniProt Gene Name	Hsd17b3
UniProt Synonym Gene Names	Edh17b3; 17-beta-HSD 3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.