

Human Nuclear Transcription Factor Y Gamma (NFYC) ELISA Kit | NFYC elisa kit

Human Nuclear Transcription Factor Y Gamma (NFYC) ELISA Kit

CATALOG #
AAA186902

REACTIVITY
Human


PRODUCT SYNONYMS

Nuclear Transcription Factor Y Gamma (NFYC); Human Nuclear Transcription Factor Y Gamma (NFYC) ELISA Kit; CBF-C; NFYg; CBFC; H1TF2A; HAP5; HSM; NF-YC; hCBF-C; CAAT box DNA-binding protein subunit C; Nuclear transcription factor Y subunit C; Transactivator HSM-1/2; NFYC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Nuclear Transcription Factor Y Gamma (NFYC). No significant cross-reactivity or interference between Nuclear Transcription Factor Y Gamma (NFYC) and analogues was observed.
Sequence Length	334
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates, Cell Lysates and Other Biological Fluids
Detection Range	0.312-20ng/mL
Sensitivity	< 0.114ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Nuclear Transcription Factor Y Gamma (NFYC) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Nuclear Transcription Factor Y Gamma (NFYC) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NFYC elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Nuclear Transcription Factor Y Gamma (NFYC). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Nuclear Transcription Factor Y Gamma (NFYC). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Nuclear Transcription Factor Y Gamma (NFYC), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Nuclear Transcription Factor Y Gamma (NFYC) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	217272835
NCBI GeneID	4802
NCBI Accession #	NP_001136062.1
NCBI GenBank Nucleotide #	NM_001142590.1
UniProt Accession #	Q13952; Q59GY4; Q5T6K8; Q5T6K9; Q5T6L1; Q5TZR6; Q92869; Q9HBX1; B4DUS6; B4DW63; D3DPV9; F8VWM3
Molecular Weight	37,066 Da
NCBI Official Full Name	nuclear transcription factor Y subunit gamma isoform 5
NCBI Official Synonym Full Names	nuclear transcription factor Y subunit gamma
NCBI Official Symbol	NFYC
NCBI Official Synonym Symbols	HSM; CBFC; HAP5; CBF-C; NF-YC; H1TF2A
NCBI Protein Information	nuclear transcription factor Y subunit gamma
UniProt Protein Name	Nuclear transcription factor Y subunit gamma
UniProt Gene Name	NFYC
UniProt Synonym Gene Names	NF-YC

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.