

Mouse Podocan (PODN) ELISA Kit | PODN elisa kit

Mouse Podocan (PODN) ELISA Kit

CATALOG #
AAA187013

REACTIVITY
Mouse


PRODUCT SYNONYMS

Podocan (PODN); Mouse Podocan (PODN) ELISA Kit; PCAN; SLRR5A; PODN elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Podocan (PODN). No significant cross-reactivity or interference between Podocan (PODN) and analogues was observed.
Sequence Length	399
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates and Other Biological Fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.057ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Podocan (PODN) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Podocan (PODN) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PODN elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Podocan (PODN). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Podocan (PODN). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Podocan (PODN), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Podocan (PODN) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	552953828
NCBI GeneID	242608
NCBI Accession #	NP_001272885.1
NCBI GenBank Nucleotide #	NM_001285956.1
UniProt Accession #	Q7TQ62; Q8BM45; A2A8F3
Molecular Weight	45,196 Da
NCBI Official Full Name	podocan
NCBI Official Synonym Full Names	podocan
NCBI Official Symbol	Podn
NCBI Official Synonym Symbols	Pcan; SLRR5A; 9430070G18
NCBI Protein Information	podocan
UniProt Protein Name	Podocan
UniProt Gene Name	Podn

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/podocan-podn/187013>