

Transcription factor p65 Recombinant Protein | NFKB3 recombinant protein

Recombinant Human Transcription factor p65 protein

CATALOG #
AAA18725

HOST
E Coli


PRODUCT SYNONYMS

Transcription factor p65; Recombinant Human Transcription factor p65 protein; Nuclear factor NF-kappa-B p65 subunit; Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3; NFKB3 recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Liquid containing glycerol
Sequence Positions	1-210aa; Partial
Sequence	MDELFLIFP AEPAQASGPY VEIIQPKQR GMRFYKCEG RSAGSIPGER STDTTKTHPT IKINGYTGP TVRISLVTKD PPHRPHPEL VGKDCRDGFY EAELCPDRCI HSFQNLGIQC VKKRDLEQAI SQRIQTNNNP FQVPIEEQRG DYDLNAVRLC FQVTVRDPSP RPLRLPPVLS HPIFDNRAPN TAEKICRVN RNSGSLGGD
Sequence Length	210
Preparation and Storage	Store at -20 degree C, for extended storage, conserve at -20 degree C or -80 degree C.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for NFKB3 recombinant protein	NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF-kappa-B p65-p65 complex appears to be involved in invasion-mediated activation of IL-8 expression. The inhibitory effect of I-kappa-B upon NF-kappa-B the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-kappa-B complex. Associates with chromatin at the NF-kappa-B promoter region via association with DDX1. Essential for cytokine gene expression in T-cells
Product Categories/Family for NFKB3 recombinant protein	Immunology

References

Isolation of a rel-related human cDNA that potentially encodes the 65-kD subunit of NF-kappa B. Ruben S.M., Dillon P.J., Schreck R., Henkel T., Chen C.-H., Maher M., Baeuerle P.A., Rosen C.A. *Science* 251:1490-1493(1991)

Genomic organization of the gene encoding the p65 subunit of NF-kappa B multiple variants of the p65 protein may be generated by alternative splicing. Deloukas P., van Loon A.P.G.M. *Hum. Mol. Genet.* 2:1895-1900(1993)

An alternatively spliced transcript, p65 delta 2, of the gene encoding the p65 subunit of the transcription factor NF-kappa B. Lyle R., Valleley E.M., Sharpe P.T., Hewitt J.E. *Gene* 138:265-266(1994)

SeattleSNPs variation discovery resource

Functional characterization of the NF-kappa B p65 transcriptional activator and an alternatively spliced derivative. Ruben S.M., Narayanan R., Klement J.F., Chen C.-H., Rosen C.A. *Mol. Cell. Biol.* 12:444-454(1992)

A novel complex between the p65 subunit of NF-kappa B and c-Rel binds to a DNA element involved in the phorbol ester induction of the human urokinase gene. Hansen S.K., Nerlov C., Zabel U., Verde P., Johnsen M., Baeuerle P.A., Blasi F. *EMBO J.* 11:205-213(1992)

I kappa B/MAD-3 masks the nuclear localization signal of NF-kappa B p65 and requires the transactivation domain to inhibit NF-kappa B p65 DNA binding. Ganchi P.A., Sun S.C., Greene W.C., Ballard D.W. *Mol. Biol. Cell* 3:1339-1352(1992)

Activation of multiple NF-kappa B/Rel DNA-binding complexes by tumor necrosis factor. Beg A.A., Baldwin A.S. Jr. *Oncogene* 9:1487-1492(1994)

Regulation of intercellular adhesion molecule-1 gene by tumor necrosis factor-alpha is mediated by the nuclear factor-kappaB heterodimers p65/p65 and p65/c-Rel in the absence of p50. Aoudjit F., Brochu N., Belanger B., Stratowa C., Hiscott J., Audette M. *Cell Growth Differ.* 8:335-342(1997)

A new member of the IkappaB protein family, IkappaB epsilon, inhibits RelA (p65)-mediated NF-kappaB transcription. Li Z., Nabel G.J. *Mol. Cell. Biol.* 17:6184-6190(1997)

IKAP is a scaffold protein of the IkappaB kinase complex. Cohen L., Henzel W.J., Baeuerle P.A. *Nature* 395:292-296(1998)

IkappaB kinases phosphorylate NF-kappaB p65 subunit on serine 536 in the transactivation domain. Sakurai H., Chiba H., Miyoshi H., Sugita T., Toriumi W.J. *Biol. Chem.* 274:30353-30356(1999)

NF-kappaB subunit p65 binds to 53BP2 and inhibits cell death induced by 53BP2. Yang J.-P., Hori M., Takahashi N., Kawabe T., Kato H., Okamoto T. *Oncogene* 18:5177-5186(1999)

Yersinia enterocolitica invasin protein triggers IL-8 production in epithelial cells via activation of Rel p65-p65 homodimers. Schulte R., Grassl G.A., Preger S., Fessele S., Jacobi C.A., Schaller M., Nelson P.J., Autenrieth I.B. *FASEB J.* 14:1471-1484(2000)

Inhibition of nuclear factor-kappaB-mediated transcription by association with the amino-terminal enhancer of split, a Groucho-related protein lacking WD40 repeats. Tetsuka T., Uranishi H., Imai H., Ono T., Sonta S., Takahashi N., Asamitsu K., Okamoto T.J. *Biol. Chem.* 275:4383-4390(2000)

Tumor necrosis factor alpha-induced phosphorylation of RelA/p65 on Ser529 is controlled by casein kinase II. Wang D., Westerheide S.D., Hanson J.L., Baldwin A.S. Jr. *J. Biol. Chem.* 275:32592-32597(2000)

The tumor suppressor protein menin interacts with NF-kappaB proteins and inhibits NF-kappaB-mediated transactivation. Heppner C., Bilimoria K.Y., Agarwal S.K., Kester M., Whitty L.J., Guru S.C., Chandrasekharappa S.C., Collins F.S., Spiegel A.M., Marx S.J., Burns A.L. *Oncogene* 20:4917-4925(2001)

Duration of nuclear NF-kappaB action regulated by reversible acetylation. Chen L.F., Fischle W., Verdin E., Greene W.C. *Science* 293:1653-1657(2001)

A novel protein overexpressed in hepatoma accelerates export of NF-kappa B from the nucleus and inhibits p53-dependent apoptosis. Higashitsuji H., Higashitsuji H., Nagao T., Nonoguchi K., Fujii S., Itoh K., Fujita J. *Cancer Cell* 2:335-346(2002)

Acetylation of RelA at discrete sites regulates distinct nuclear functions of NF-kappaB. Chen L.F., Mu Y., Greene W.C. *EMBO J.* 21:6539-6548(2002)

The phosphorylation status of nuclear NF-kappa B determines its association with CBP/p300 or HDAC-1. Zhong H., May M.J., Jimi E., Ghosh S. *Mol. Cell* 9:625-636(2002)

Transcriptional activation of the NF-kappaB p65 subunit by mitogen- and stress-activated protein kinase-1 (MSK1). Vermeulen L., De Wilde G., Van Damme P., Vanden Berghe W., Haegeman G. *EMBO J.* 22:1313-1324(2003)

Post-activation turn-off of NF-kappa B-dependent transcription is regulated by acetylation of p65. Kiernan R., Bres V., Ng R.W., Coudart M.-P., El Messaoudi S., Sartet C., Jin D.-Y., Emiliani S., Benkirane M.J. *Biol. Chem.* 278:2758-2766(2003)

RING finger protein A07 supports NF-kappaB-mediated transcription by interacting with the transactivation domain of the p65 subunit. Asamitsu K., Tetsuka T., Kanazawa S., Okamoto T.J. *Biol. Chem.* 278:26879-26887(2003)

Glycogen synthase kinase-3 beta regulates NF-kappa B1/p105 stability. Demarchi F., Bertoli C., Sandy P., Schneider C.J. *Biol. Chem.* 278:39583-39590(2003)

Regulation of NF-kappaB signaling by Pin1-dependent prolyl isomerization and ubiquitin-mediated proteolysis of p65/RelA. Ryo A., Suizu F., Yoshida Y., Perrem K., Liou Y.C., Wulf G., Rottapel R., Yamaoka S., Lu K.P. *Mol. Cell* 12:1413-1426(2003)

Identification of a ZU5 and death domain-containing inhibitor of NF-kappaB. Zhang J., Xu L.-G., Han K.-J., Shu H.-B.J. *Biol. Chem.* 279:17819-17825(2004)

Suppression of MEK/ERK signaling pathway enhances cisplatin-induced NF-kappaB activation by protein phosphatase 4-mediated NF-kappaB p65 Thr dephosphorylation. Yeh P.Y., Yeh K.H., Chuang S.E., Song Y.C., Cheng A.L.J. *Biol. Chem.* 279:26143-26148(2004)

Degradation of promoter-bound p65/RelA is essential for the prompt termination of the nuclear factor kappaB response. Saccani S., Marazzi I., Beg A.A., Natoli G.J. *Exp. Med.* 200:107-113(2004)

Identification of beta-arrestin2 as a G protein-coupled receptor-stimulated regulator of NF-kappaB pathways. Gao H., Sun Y., Wu Y., Luan B., Wang Y., Qu B., Pei G. *Mol. Cell* 14:303-317(2004)

The candidate tumour suppressor protein ING4 regulates brain tumour growth and angiogenesis. Garkavtsev I., Kozin S.V., Chernova O., Xu L., Winkler F., Brown E., Barnett G.H., Jain R.K. *Nature* 428:328-332(2004)

Regulation of NF-kappaB and p53 through activation of ATR and Chk1 by the ARF tumour suppressor. Rocha S., Garrett M.D., Campbell K.J., Schumm K., Perkins N.D. *EMBO J.* 24:1157-1169(2005)

IKKbeta phosphorylates p65 at S468 in transactivation domain 2. Schwabe R.F., Sakurai H. *FASEB J.* 19:1758-1760(2005)

cis-acting, element-specific transcriptional activity of differentially phosphorylated nuclear factor-kappa B. Anrather J., Racchumi G., Iadecola C.J. *Biol. Chem.* 280:244-252(2005)

COMMD proteins, a novel family of structural and functional homologs of MURR1. Burstein E., Hoberg J.E., Wilkinson A.S., Rumble J.M., Csomos R.A., Komarck C.M., Maine G.N., Wilkinson J.C., Mayo M.W., Duckett C.S.J. *Biol. Chem.* 280:22222-22232(2005)

NF-kappaB RelA phosphorylation regulates RelA acetylation. Chen L.F., Williams S.A., Mu Y., Nakano H., Duerr J.M., Buckbinder L., Greene W.C. *Mol. Cell. Biol.* 25:7966-7975(2005)

Foxp3 interacts with nuclear factor of activated T cells and NF-kappa B to repress cytokine gene expression and effector functions of T helper cells. Bettelli E., Dastrange M., Oukka M. *Proc. Natl. Acad. Sci. U.S.A.* 102:5138-5143(2005)

Respiratory syncytial virus M2-1 protein induces the activation of nuclear factor kappa B. Reimers K., Buchholz K., Werchau H. *Virology* 331:260-268(2005)

NCBI AND UNIPROT INFORMATION

NCBI GI #	223468681
NCBI GeneID	5970
NCBI Accession #	NP_001138610.1
NCBI GenBank Nucleotide #	NM_001145138.1
UniProt Accession #	Q04206; Q6GTV1; Q6SLK1
Molecular Weight	27.7kD
NCBI Official Full Name	transcription factor p65 isoform 2
NCBI Official Synonym Full Names	v-rel avian reticuloendotheliosis viral oncogene homolog A
NCBI Official Symbol	RELA
NCBI Official Synonym Symbols	p65; NFkB3
NCBI Protein Information	transcription factor p65
UniProt Protein Name	Transcription factor p65
UniProt Gene Name	RELA
UniProt Synonym Gene Names	NFkB3
UniProt Entry Name	TF65_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rela-recombinant-protein-e-coli/18725>