

Rat Proopiomelanocortin (POMC) ELISA Kit | POMC elisa kit

Rat Proopiomelanocortin (POMC) ELISA Kit

CATALOG #
AAA188153

REACTIVITY
Rat


PRODUCT SYNONYMS

Proopiomelanocortin (POMC); Rat Proopiomelanocortin (POMC) ELISA Kit; POC; ProOMC; Pro-OMC; Pro Opiomelanocortin; Corticotropin-lipotropin; POMC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Proopiomelanocortin (POMC). No significant cross-reactivity or interference between Proopiomelanocortin (POMC) and analogues was observed.
Sequence Length	235
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates and Other Biological Fluids
Detection Range	78-5,000pg/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 33pg/mL
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Proopiomelanocortin (POMC) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Proopiomelanocortin (POMC) were tested on 3 different plates, 8 replicates in each plate. $CV(\%) = SD/mean \times 100$ Intra-Assay: $CV < 10\%$ Inter-Assay: $CV < 12\%$
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for POMC elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Proopiomelanocortin (POMC). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Proopiomelanocortin (POMC). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Proopiomelanocortin (POMC), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Proopiomelanocortin (POMC) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	116889
NCBI GeneID	24664
UniProt Accession #	P01194
NCBI Official Full Name	Pro-opiomelanocortin
NCBI Official Synonym Full Names	proopiomelanocortin
NCBI Official Symbol	Pomc
NCBI Official Synonym Symbols	Pomc1; Pomc2; alphaMSH
NCBI Protein Information	pro-opiomelanocortin
UniProt Protein Name	Pro-opiomelanocortin
UniProt Gene Name	Pomc
UniProt Synonym Gene Names	Pomc2; POMC; ACTH; CLIP
UniProt Entry Name	COLI_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.