

Mouse Loricrin (LOR) ELISA Kit | LOR elisa kit

Mouse Loricrin (LOR) ELISA Kit

CATALOG #
AAA188193

REACTIVITY
Mouse


PRODUCT SYNONYMS

Loricrin (LOR); Mouse Loricrin (LOR) ELISA Kit; LOR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Loricrin (LOR). No significant cross-reactivity or interference between Loricrin (LOR) and analogues was observed.
Sequence Length	312
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates, Cell Lysates and Other Biological Fluids
Detection Range	0.156-10ng/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 0.057ng/mL
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Loricrin (LOR) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Loricrin (LOR) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for LOR elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Loricrin (LOR). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Loricrin (LOR). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Loricrin (LOR), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Loricrin (LOR) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	109255251
NCBI GeneID	4014
NCBI GenBank Nucleotide #	NP_000418.2
UniProt Accession #	P23490
NCBI Official Full Name	loricrin
NCBI Official Synonym Full Names	loricrin cornified envelope precursor protein
NCBI Official Symbol	LORICRIN
NCBI Official Synonym Symbols	LOR
NCBI Protein Information	loricrin
UniProt Protein Name	Loricrin
UniProt Gene Name	LOR
UniProt Synonym Gene Names	LRN
UniProt Entry Name	LORI_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/loricrin-lor/188193>