

Human Syntenin 1 (ST1) ELISA Kit | ST1 elisa kit

Human Syntenin 1 (ST1) ELISA Kit

CATALOG #
AAA188198

REACTIVITY
Human


PRODUCT SYNONYMS

Syntenin 1 (ST1); Human Syntenin 1 (ST1) ELISA Kit; SDCBP; MDA9; SYCL; TACIP18; Syndecan Binding Protein; Scaffold protein Pbp1; Melanoma differentiation-associated protein 9; Pro-TGF-alpha cytoplasmic domain-interacting 18; ST1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of ST1. No significant cross-reactivity or interference between ST1 and analogues was observed.
Sequence Length	298
Assay Type	Quantitative Sandwich
Samples	Human tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.062ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ST1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ST1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ST1 elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to ST1. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to ST1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ST1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of ST1 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of ST1 in human tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	20455281
NCBI GeneID	6386
UniProt Accession #	O00560
NCBI Official Full Name	Syntenin-1
NCBI Official Synonym Full Names	syndecan binding protein
NCBI Official Symbol	SDCBP
NCBI Official Synonym Symbols	ST1; MDA9; SYCL; MDA-9; TACIP18
NCBI Protein Information	syntenin-1
UniProt Protein Name	Syntenin-1
UniProt Gene Name	SDCBP
UniProt Synonym Gene Names	MDA9; SYCL; MDA-9; TACIP18
UniProt Entry Name	SDCB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.