

Rat Hexokinase 2 (HK2) ELISA Kit | HK2 elisa kit

Rat Hexokinase 2 (HK2) ELISA Kit

CATALOG #
AAA188219

REACTIVITY
Rat


PRODUCT SYNONYMS

Hexokinase 2 (HK2); Rat Hexokinase 2 (HK2) ELISA Kit; HK II; Muscle form hexokinase; HK2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Hexokinase 2 (HK2). No significant cross-reactivity or interference between Hexokinase 2 (HK2) and analogues was observed.
Sequence Length	917
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates and Other Biological Fluids
Detection Range	15.6-1,000pg/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 6.0pg/mL
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Hexokinase 2 (HK2) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Hexokinase 2 (HK2) were tested on 3 different plates, 8 replicates in each plate. $CV(\%) = SD/mean \times 100$ Intra-Assay: $CV < 10\%$ Inter-Assay: $CV < 12\%$
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HK2 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Hexokinase 2 (HK2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Hexokinase 2 (HK2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Hexokinase 2 (HK2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Hexokinase 2 (HK2) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	39963174
NCBI GeneID	3099
NCBI Official Full Name	Hexokinase 2
NCBI Official Synonym Full Names	hexokinase 2
NCBI Official Symbol	HK2
NCBI Official Synonym Symbols	HKII; HXK2
NCBI Protein Information	hexokinase-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/hexokinase-2-hk2/188219>