

Human Alcohol Dehydrogenase 4 (ADH4) ELISA Kit | ADH4 elisa kit

Human Alcohol Dehydrogenase 4 (ADH4) ELISA Kit

CATALOG #
AAA188249

REACTIVITY
Human


PRODUCT SYNONYMS

Alcohol Dehydrogenase 4 (ADH4); Human Alcohol Dehydrogenase 4 (ADH4) ELISA Kit; Alcohol dehydrogenase class II pi chain; ADH4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Alcohol Dehydrogenase 4 (ADH4). No significant cross-reactivity or interference between Alcohol Dehydrogenase 4 (ADH4) and analogues was observed.
Sequence Length	380
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates, Cell Lysates and Other Biological Fluids
Detection Range	0.78-50ng/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 0.29ng/mL
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Alcohol Dehydrogenase 4 (ADH4) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Alcohol Dehydrogenase 4 (ADH4) were tested on 3 different plates, 8 replicates in each plate. $CV(\%) = SD/mean \times 100$ Intra-Assay: $CV < 10\%$ Inter-Assay: $CV < 12\%$
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ADH4 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Alcohol Dehydrogenase 4 (ADH4). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Alcohol Dehydrogenase 4 (ADH4). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Alcohol Dehydrogenase 4 (ADH4), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Alcohol Dehydrogenase 4 (ADH4) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	71565152
NCBI GeneID	127
NCBI Accession #	NP_000661.2
NCBI GenBank Nucleotide #	NP_000661.2
UniProt Accession #	P08319
NCBI Official Full Name	all-trans-retinol dehydrogenase
NCBI Official Synonym Full Names	alcohol dehydrogenase 4 (class II), pi polypeptide
NCBI Official Symbol	ADH4
NCBI Official Synonym Symbols	ADH-2; HEL-S-4
NCBI Protein Information	all-trans-retinol dehydrogenase [NAD(+)] ADH4
UniProt Protein Name	Alcohol dehydrogenase 4
UniProt Gene Name	ADH4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/alcohol-dehydrogenase-4-adh4/188249>