

Rat Stromal Cell Derived Factor 1 (SDF1) ELISA Kit | SDF1 elisa kit

Rat Stromal Cell Derived Factor 1 (SDF1) High Sensitive ELISA Kit

CATALOG # AAA188482	REACTIVITY Rat	APPLICATIONS ELISA
-------------------------------	--------------------------	------------------------------


PRODUCT SYNONYMS

Stromal Cell Derived Factor 1 (SDF1); Rat Stromal Cell Derived Factor 1 (SDF1) High Sensitive ELISA Kit; Stromal Cell Derived Factor 1; CXCL12; PBSF; P-BSF; SCYB12; SDF1-A; SDF1-B; SDF1A; SDF1B; TLSF-A; TLSF-B; TPAR1; Pre-B Cell Growth-Stimulating Factor; Chemokine C-X-C-Motif Ligand 12; SDF1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of High Sensitive Stromal Cell Derived Factor 1 (SDF1). No significant cross-reactivity or interference between High Sensitive Stromal Cell Derived Factor 1 (SDF1) and analogues was observed.
Sequence Length	89
Assay Type	Double-antibody Sandwich
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	15.6-1,000pg/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 6.2pg/mL
Applicable Applications for SDF1 elisa kit	ELISA
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level High Sensitive Stromal Cell Derived Factor 1 (SDF1) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level High Sensitive Stromal Cell Derived Factor 1 (SDF1) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to High Sensitive Stromal Cell Derived Factor 1 (SDF1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to High Sensitive Stromal Cell Derived Factor 1 (SDF1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain High Sensitive Stromal Cell Derived Factor 1 (SDF1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of High Sensitive Stromal Cell Derived Factor 1 (SDF1) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	11559933
NCBI GeneID	24772
NCBI Accession #	NP_071513.1
NCBI GenBank Nucleotide #	NP_071513.1
UniProt Accession #	Q9QZD1
NCBI Official Full Name	stromal cell-derived factor 1 isoform alpha
NCBI Official Synonym Full Names	C-X-C motif chemokine ligand 12
NCBI Official Symbol	Cxcl12
NCBI Official Synonym Symbols	Sdf1
NCBI Protein Information	stromal cell-derived factor 1
UniProt Protein Name	Chemokine (C-X-C motif) ligand 12 (Stromal cell-derived factor 1)
UniProt Gene Name	Cxcl12
UniProt Entry Name	Q9QZD1_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/stromal-cell-derived-factor-1-sdf1/188482>