

Human Transglutaminase 2 (TGM2) ELISA Kit | TGM2 elisa kit

Human Transglutaminase 2 (TGM2) High Sensitive ELISA Kit

CATALOG # AAA188554	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

Transglutaminase 2 (TGM2); Human Transglutaminase 2 (TGM2) High Sensitive ELISA Kit; Transglutaminase 2; tTG; TG2; C Polypeptide; Protein-Glutamine-Gamma-Glutamyltransferase; Tissue; Transglutaminase C; Transglutaminase H; TGM2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of High Sensitive Transglutaminase 2 (TGM2). No significant cross-reactivity or interference between High Sensitive Transglutaminase 2 (TGM2) and analogues was observed.
Sequence Length	687
Assay Type	Double-antibody Sandwich
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 0.060ng/mL
Applicable Applications for TGM2 elisa kit	ELISA
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level High Sensitive Transglutaminase 2 (TGM2) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level High Sensitive Transglutaminase 2 (TGM2) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to High Sensitive Transglutaminase 2 (TGM2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to High Sensitive Transglutaminase 2 (TGM2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain High Sensitive Transglutaminase 2 (TGM2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of High Sensitive Transglutaminase 2 (TGM2) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	39777597
NCBI GeneID	7052
NCBI Accession #	NP_004604.2
NCBI GenBank Nucleotide #	NP_004604.2
UniProt Accession #	P21980
NCBI Official Full Name	protein-glutamine gamma-glutamyltransferase 2 isoform a
NCBI Official Synonym Full Names	transglutaminase 2
NCBI Official Symbol	TGM2
NCBI Official Synonym Symbols	TGC; TG(C)
NCBI Protein Information	protein-glutamine gamma-glutamyltransferase 2
UniProt Protein Name	Protein-glutamine gamma-glutamyltransferase 2
UniProt Gene Name	TGM2
UniProt Synonym Gene Names	TG(C); TGC; TGase C; TGase H; TGase-2
UniProt Entry Name	TGM2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.