

Human Dopa Decarboxylase (DDC) ELISA Kit | DDC elisa kit

Human Dopa Decarboxylase (DDC) High Sensitive ELISA Kit

CATALOG # AAA188595	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

Dopa Decarboxylase (DDC); Human Dopa Decarboxylase (DDC) High Sensitive ELISA Kit; Dopa Decarboxylase; AADC; Aromatic L-Amino Acid Decarboxylase; DOPA decarboxylase; DDC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of High Sensitive Dopa Decarboxylase (DDC). No significant cross-reactivity or interference between High Sensitive Dopa Decarboxylase (DDC) and analogues was observed.
Sequence Length	480
Assay Type	Double-antibody Sandwich
Samples	tissue homogenates, cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 0.056ng/mL
Applicable Applications for DDC elisa kit	ELISA
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level High Sensitive Dopa Decarboxylase (DDC) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level High Sensitive Dopa Decarboxylase (DDC) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to High Sensitive Dopa Decarboxylase (DDC). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to High Sensitive Dopa Decarboxylase (DDC). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain High Sensitive Dopa Decarboxylase (DDC), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of High Sensitive Dopa Decarboxylase (DDC) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	311033369
NCBI GeneID	1644
UniProt Accession #	P20711
NCBI Official Full Name	Aromatic-L-amino-acid decarboxylase
NCBI Official Synonym Full Names	dopa decarboxylase
NCBI Official Symbol	DDC
NCBI Official Synonym Symbols	AADC
NCBI Protein Information	aromatic-L-amino-acid decarboxylase
UniProt Protein Name	Aromatic-L-amino-acid decarboxylase
UniProt Gene Name	DDC
UniProt Synonym Gene Names	AADC; AADC; DDC
UniProt Entry Name	DDC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/dopa-decarboxylase-ddc/188595>