

Mouse Fibrinogen Beta Chain (FGB) ELISA Kit | FGB elisa kit

Mouse Fibrinogen Beta Chain (FGB) Mini Samples ELISA Kit

CATALOG # AAA188733	REACTIVITY Mouse	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

Fibrinogen Beta Chain (FGB); Mouse Fibrinogen Beta Chain (FGB) Mini Samples ELISA Kit; Fibrinogen Beta Chain; FG-B; Fibrinopeptide B; Fibrinogen Beta; FGB elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Mini Samples Fibrinogen Beta Chain (FGB). No significant cross-reactivity or interference between Mini Samples Fibrinogen Beta Chain (FGB) and analogues was observed.
Sequence Length	491
Assay Type	Double-antibody Sandwich
Samples	Plasma
Detection Range	15.6-1,000ng/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 7.1ng/mL
Applicable Applications for FGB elisa kit	ELISA
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Mini Samples Fibrinogen Beta Chain (FGB) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Mini Samples Fibrinogen Beta Chain (FGB) were tested on 3 different plates, 8 replicates in each plate. $CV(\%) = SD/mean \times 100$ Intra-Assay: $CV < 10\%$ Inter-Assay: $CV < 12\%$
Test Principle	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Mini Samples Fibrinogen Beta Chain (FGB). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Mini Samples Fibrinogen Beta Chain (FGB). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Mini Samples Fibrinogen Beta Chain (FGB), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of Mini Samples Fibrinogen Beta Chain (FGB) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	7924018
NCBI GeneID	2244
NCBI Official Full Name	fibrinogen beta chain
NCBI Official Synonym Full Names	fibrinogen beta chain
NCBI Official Symbol	FGB
NCBI Official Synonym Symbols	HEL-S-78p
NCBI Protein Information	fibrinogen beta chain

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.