

Rat Laminin (LN) CLIA Kit | LN clia kit

Rat Laminin (LN) CLIA Kit

CATALOG # AAA188832	REACTIVITY Rat	APPLICATIONS CLIA (Chemiluminescence Immunoassay)
-------------------------------	--------------------------	---


PRODUCT SYNONYMS

Laminin (LN); Rat Laminin (LN) CLIA Kit; Laminin; LAM; LN clia kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Laminin (LN). No significant cross-reactivity or interference between Laminin (LN) and analogues was observed.
Sequence Length	1111
Assay Type	Double-antibody Sandwich
Samples	Serum, plasma, tissue homogenates and other biological fluids
Detection Range	0.27-200ng/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 0.10ng/mL
Applicable Applications for LN clia kit	CLIA (Chemiluminescence Immunoassay)
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Laminin (LN) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Laminin (LN) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The microplate provided in this kit has been pre-coated with an antibody specific to Laminin (LN). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Laminin (LN). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Laminin (LN) level in the sample or standard.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	34232
NCBI GeneID	3918
NCBI Official Full Name	Laminin
NCBI Official Synonym Full Names	laminin subunit gamma 2
NCBI Official Symbol	LAMC2
NCBI Official Synonym Symbols	B2T; CSF; EBR2; BM600; EBR2A; LAMB2T; LAMNB2
NCBI Protein Information	laminin subunit gamma-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-clia-kits/laminin-ln/188832>