

Mouse Osteoprotegerin (OPG) CLIA Kit | OPG clia kit

Mouse Osteoprotegerin (OPG) CLIA Kit

CATALOG # AAA188859	REACTIVITY Mouse	APPLICATIONS CLIA (Chemiluminescence Immunoassay)
-------------------------------	----------------------------	---


PRODUCT SYNONYMS

Osteoprotegerin (OPG); Mouse Osteoprotegerin (OPG) CLIA Kit; Osteoprotegerin; TNFRSF11B; TNFRSF11-B; OCIF; OPG; TR1; Tumor Necrosis Factor Receptor Superfamily Member 11b; Osteoclastogenesis Inhibitory Factor; OPG clia kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Osteoprotegerin (OPG). No significant cross-reactivity or interference between Osteoprotegerin (OPG) and analogues was observed.
Sequence Length	401
Assay Type	Double-antibody Sandwich
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	27.4-20,000pg/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 11.8pg/mL
Applicable Applications for OPG clia kit	CLIA (Chemiluminescence Immunoassay)
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Osteoprotegerin (OPG) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Osteoprotegerin (OPG) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The microplate provided in this kit has been pre-coated with an antibody specific to Osteoprotegerin (OPG). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Osteoprotegerin (OPG). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Osteoprotegerin (OPG) level in the sample or standard.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	113930716
NCBI GeneID	18383
NCBI Accession #	NP_032790.3
NCBI GenBank Nucleotide #	NP_032790.3
UniProt Accession #	O08712
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 11B
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily, member 11b (osteoprotegerin)
NCBI Official Symbol	Tnfrsf11b
NCBI Official Synonym Symbols	Opg; TR1; OCIF
NCBI Protein Information	tumor necrosis factor receptor superfamily member 11B
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 11B
UniProt Gene Name	Tnfrsf11b
UniProt Synonym Gene Names	Ocif; Opg

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-clia-kits/osteoprotegerin-opg/188859>