

Human Tumor Necrosis Factor Alpha (TNFa) CLIA Kit | TNFa clia kit

Human Tumor Necrosis Factor Alpha (TNFa) CLIA Kit

CATALOG # AAA188874	REACTIVITY Human	APPLICATIONS CLIA (Chemiluminescence Immunoassay)
-------------------------------	----------------------------	---


PRODUCT SYNONYMS

Tumor Necrosis Factor Alpha (TNFa); Human Tumor Necrosis Factor Alpha (TNFa) CLIA Kit; Tumor Necrosis Factor Alpha; DIF; TNF-A; TNFSF2; Cachectin; Tumor Necrosis Factor Ligand Superfamily Member 2; TNFa clia kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Tumor Necrosis Factor Alpha (TNFa). No significant cross-reactivity or interference between Tumor Necrosis Factor Alpha (TNFa) and analogues was observed.
Sequence Length	233
Assay Type	Double-antibody Sandwich
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	0.14-100pg/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 0.05pg/mL
Applicable Applications for TNFa clia kit	CLIA (Chemiluminescence Immunoassay)
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Tumor Necrosis Factor Alpha (TNFa) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Tumor Necrosis Factor Alpha (TNFa) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The microplate provided in this kit has been pre-coated with an antibody specific to Tumor Necrosis Factor Alpha (TNFa). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Tumor Necrosis Factor Alpha (TNFa). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Tumor Necrosis Factor Alpha (TNFa) level in the sample or standard.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	25952111
NCBI GeneID	7124
NCBI Accession #	NP_000585.2
NCBI GenBank Nucleotide #	NP_000585.2
UniProt Accession #	P01375
NCBI Official Full Name	tumor necrosis factor
NCBI Official Synonym Full Names	tumor necrosis factor
NCBI Official Symbol	TNF
NCBI Official Synonym Symbols	DIF; TNFA; TNFSF2; TNLG1F; TNF-alpha
NCBI Protein Information	tumor necrosis factor
UniProt Protein Name	Tumor necrosis factor
UniProt Gene Name	TNF
UniProt Synonym Gene Names	TNFA; TNFSF2; TNF-a; NTF; ICD1; ICD2
UniProt Entry Name	TNFA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-clia-kits/tumor-necrosis-factor-alpha-tnfa/188874>