

Human Apoptosis Inducing Factor (AIF) CLIA Kit | AIF clia kit

Human Apoptosis Inducing Factor (AIF) CLIA Kit

CATALOG # AAA189003	REACTIVITY Human	APPLICATIONS CLIA (Chemiluminescence Immunoassay)
-------------------------------	----------------------------	---


PRODUCT SYNONYMS

Apoptosis Inducing Factor (AIF); Human Apoptosis Inducing Factor (AIF) CLIA Kit; Apoptosis Inducing Factor; PDCD8; Programmed Cell Death protein 8; Apoptosis-inducing factor 1; mitochondrial; AIF clia kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Apoptosis Inducing Factor (AIF). No significant cross-reactivity or interference between Apoptosis Inducing Factor (AIF) and analogues was observed.
Sequence Length	261
Assay Type	Double-antibody Sandwich
Samples	Serum, plasma, tissue homogenates and other biological fluids
Detection Range	2.74-2,000pg/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 1.17pg/mL
Applicable Applications for AIF clia kit	CLIA (Chemiluminescence Immunoassay)
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Apoptosis Inducing Factor (AIF) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Apoptosis Inducing Factor (AIF) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The microplate provided in this kit has been pre-coated with an antibody specific to Apoptosis Inducing Factor (AIF). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Apoptosis Inducing Factor (AIF). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Apoptosis Inducing Factor (AIF) level in the sample or standard.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	195927004
NCBI Accession #	NP_001124318.1
NCBI GenBank Nucleotide #	NP_001124318.1
UniProt Accession #	O95831
NCBI Official Full Name	apoptosis-inducing factor 1, mitochondrial isoform 4
UniProt Protein Name	Apoptosis-inducing factor 1, mitochondrial
UniProt Gene Name	AIFM1
UniProt Synonym Gene Names	AIF; PDCD8
UniProt Entry Name	AIFM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.